

* Linear Algebra [3M]

Topics :-

- Determinants
- Inverse of a matrix
- Rank of matrix
- Homogeneous & non-homogeneous linear eqⁿ
- Eigen values & Eigen vectors
- Cayley - Hamilton theorem

Textbook :-

Matrices by A.R. Nasistha

* Determinants *

* Properties of Determinants :-

Point I : Value of the determinant will not be change if rows and columns are interchange.

$$\text{i.e., } |A| = |A^T|$$

Ex.

$$|A| = \begin{vmatrix} 5 & 4 \\ 1 & 2 \end{vmatrix} = 10 - 4 = 6$$

$$|A^T| = \begin{vmatrix} 5 & 1 \\ 4 & 2 \end{vmatrix} = 10 - 4 = 6$$

$$\therefore |A| = |A^T|$$

Point II : Value of the determinant is multiplied by -1 if two rows & two columns are interchange.

Ex.

$$|A| = \begin{vmatrix} 1 & 2 \\ 3 & 4 \end{vmatrix} = 4 - 6 = -2$$

$$|A| = \begin{vmatrix} 3 & 4 \\ 1 & 2 \end{vmatrix} = 6 - 4 = 2$$

Point III : Value of the determinant can be zero in the following cases :-

i) The elements in two rows or two columns are identical.

Ex.

$$|A| = \begin{vmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 1 & 2 & 3 \end{vmatrix} = 0$$

same

ii) The elements in two rows or two columns are proportional to each other.

Ex.

$$|A| = \begin{vmatrix} 1 & 2 & 3 \\ 0 & 1 & 5 \\ 3 & 6 & 9 \end{vmatrix} = 0$$

proportional

iii) All elements in a row or column are zeros.

Ex.

$$|A| = \begin{vmatrix} 1 & 2 & 3 \\ 0 & 0 & 0 \\ 4 & 7 & 3 \end{vmatrix} = 0$$

iv) The elements in a determinant are of consecutive order (continuous order)

→ valid for ~~at~~ 3×3 & high order matrices

Ex.

$$|A| = \begin{vmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{vmatrix} = 0$$

we can start 1st element from any no.

Point IV: The determinant of upper triangular, lower triangular, diagonal, scalar & Identity matrix is the product of its diagonal elements.

1) Upper triangular matrix

Ex.

$$|A| = \begin{vmatrix} 1 & 2 & 3 \\ 0 & -4 & 2 \\ 0 & 0 & 7 \end{vmatrix} = 1 \times (-4) \times 7 = -28$$

2) Lower triangular matrix

Ex.

$$|A| = \begin{vmatrix} 1 & 0 & 0 \\ 2 & 3 & 0 \\ 4 & 5 & 7 \end{vmatrix} = 1 \times 3 \times 7 = 21$$

3) Diagonal matrix

Ex.

$$|A| = \begin{vmatrix} 3 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 5 \end{vmatrix} = 3 \times 2 \times 5 = 30$$

A matrix is diagonal iff at least one diagonal element should be non zero & all other non-diagonal elements should be zero.

4) Scalar matrix

Ex.

$$|A| = \begin{vmatrix} 3 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 3 \end{vmatrix} = 3 \times 3 \times 3 = 27$$

→ Diagonal elements should be same

→ Non-diagonal elements should be zero.

5) Identity matrix

Ex.

$$|I| = \begin{vmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{vmatrix} = 1 \times 1 \times 1 = 1$$

→ Determinant of Identity matrix is always 1.

Point V: If A is $N \times N$ matrix then

$$|KA| = K^n |A|$$

where n = order of matrix A

Ex.

$$|A| = \begin{vmatrix} 1 & 2 \\ 3 & 4 \end{vmatrix} = 4 - 6 = -2$$

$$|3A| = 3^2 \cdot |A| = 9 \times (-2) = -18$$

Point VI: If each element of a row or column contains sum of 2 elements then the determinant can be expressed as sum of two determinants of same order.

Ex.

$$|A| = \begin{vmatrix} 1 & 1^2 & 1^3+1 \\ 2 & 2^2 & 2^3+4 \\ 3 & 3^2 & 3^3+5 \end{vmatrix}$$


$$= \begin{vmatrix} 1 & 1^2 & 1^3 \\ 2 & 2^2 & 2^3 \\ 3 & 3^2 & 3^3 \end{vmatrix} + \begin{vmatrix} 1 & 1^2 & 1 \\ 2 & 2^2 & 4 \\ 3 & 3^2 & 5 \end{vmatrix}$$

* Note 1: Consider the matrix

$$A = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}$$

Method I:

$$|A| = a_{11}(a_{22}a_{33} - a_{23}a_{32}) - a_{12}(a_{21}a_{33} - a_{23}a_{31}) + a_{13}(a_{21}a_{32} - a_{22}a_{31})$$

Method II:

$$|A| = a_{13}a_{21}a_{32} + a_{11}a_{22}a_{33} + a_{12}a_{23}a_{31} - a_{33}a_{21}a_{12} - a_{31}a_{22}a_{13} - a_{32}a_{23}a_{11}$$

→ If determinant contains more no. of zeros use method I.

Ex. 1) Find determinant of

$$A = \begin{pmatrix} 1 & 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 & 1 \end{pmatrix}$$

$$\begin{vmatrix} 2 & 1 & 2 & 2 & 1 \\ & 2 & 1 & 2 & \\ 1 & 2 & 2 & 1 & 2 \end{vmatrix}$$

$$= 8 + 1 + 8 - 4 - 4 - 4$$

$$= 5$$

Ex: 2) $A = \begin{pmatrix} 5 & 0 & 2 \\ 0 & 3 & 0 \\ 2 & 0 & 1 \end{pmatrix}$

$$= 3(5 - 4) = 3$$

Ques

Ex. 3) The following represents eqⁿ of straight line

$$\begin{vmatrix} x & 2 & 4 \\ y & 8 & 0 \\ 1 & 1 & 1 \end{vmatrix} = 0$$

The line passes through

a) (0,0) ☒ b) (3,4) c) (4,3) d) (4,4)

$$x(8) - 2(y) + 4(y - 8)$$

$$8x - 2y + 4y - 32 = 0$$

$$8x + 2y = 32$$

$$4x + y = 16$$

$$\therefore 4(3) + y = 16$$

$$x = 3 \quad \& \quad y = 4$$

Note 2:- Consider the matrix

$$A = \begin{pmatrix} + & - & + & - \\ a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{pmatrix}_{4 \times 4}$$

Method I :- (Complicated)

$$|A| = a_{11} \begin{vmatrix} a_{22} & a_{23} & a_{24} \\ a_{32} & a_{33} & a_{34} \\ a_{42} & a_{43} & a_{44} \end{vmatrix} - a_{12} \begin{vmatrix} a_{21} & a_{23} & a_{24} \\ a_{31} & a_{33} & a_{34} \\ a_{41} & a_{43} & a_{44} \end{vmatrix}$$

$$+ a_{13} \begin{vmatrix} a_{21} & a_{22} & a_{24} \\ a_{31} & a_{32} & a_{34} \\ a_{41} & a_{42} & a_{44} \end{vmatrix} - a_{14} \begin{vmatrix} a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \\ a_{41} & a_{42} & a_{43} \end{vmatrix}$$

Method II :- (Preferable)

Ex

$$A = \begin{pmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{pmatrix}_{4 \times 4}$$

Procedure:

Step 1 : Among the given element of a matrix, select any non zero element.

Step 2 : Make all elements above & below or left & right of the selected element as zero using row & column operations.

Therefore, the matrix is

$$A = \begin{pmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ 0 & b_{22} & b_{23} & b_{24} \\ 0 & b_{32} & b_{33} & b_{34} \\ 0 & b_{42} & b_{43} & b_{44} \end{pmatrix}$$

→ 1st column has more no. of zeros so we take the determinant along 1st column.

$$|A| = a_{11} \begin{vmatrix} b_{22} & b_{23} & b_{24} \\ b_{32} & b_{33} & b_{34} \\ b_{42} & b_{43} & b_{44} \end{vmatrix}$$

Ex. 1) Find the determinant of

$$A = \begin{pmatrix} 1 & -1 & 0 & 1 \\ 1 & 2 & 3 & 4 \\ 2 & 3 & 4 & 9 \\ 0 & 1 & 0 & 2 \end{pmatrix}$$

$$R_2 \rightarrow R_2 - R_1$$

$$R_3 \rightarrow R_3 - 2R_1$$

$$A = \begin{pmatrix} 1 & -1 & 0 & 1 \\ 0 & 3 & 3 & 3 \\ 0 & 5 & 4 & 7 \\ 0 & 1 & 0 & 2 \end{pmatrix}$$

$$|A| = \begin{vmatrix} 1 & 3 & 3 & 3 \\ 0 & 5 & 4 & 7 \\ 0 & 1 & 0 & 2 \end{vmatrix}$$

$$1 [21 - 12] + 2 [12 - 15]$$

$$9 + 2(-3)$$

$$9 - 6 = 3$$

Problem 1 :- If A has m rows & $m+5$ columns & B has n rows and $11-n$ columns. The orders of A and B if AB and BA are defined?

$$(A) \quad m \times (m+5) \quad (B) \quad n \times (11-n)$$

$$(B) \quad n \times (11-n) \quad (A) \quad m \times (m+5)$$

$$\therefore m+5 = n$$

$$11-n = m$$

$$m - n = -5$$

$$m + n = 11$$

$$2m = 6$$

$$m = 3$$

$$n = 8$$

Therefore, orders are $A_{(3,8)}$ & $B_{(8,3)}$ resp

Problem 2 :- If $A = (a_{ij})_{m \times n}$ such that $a_{ij} = i+j, \forall i, j$ then sum of all element of A is?

$$A = \begin{pmatrix} 1+1 & 1+2 & 1+3 & \dots & 1+n \\ 2+1 & 2+2 & 2+3 & \dots & 2+n \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ m+1 & m+2 & m+3 & \dots & m+n \end{pmatrix}$$

according to point VI

$$A = \begin{pmatrix} 1 & 1 & \dots & 1 \\ 2 & 2 & \dots & 2 \\ 3 & 3 & \dots & 3 \\ \vdots & \vdots & \ddots & \vdots \\ m & m & \dots & m \end{pmatrix} + \begin{pmatrix} 1 & 2 & 3 & \dots & n \\ 1 & 2 & 3 & \dots & n \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & 2 & 3 & \dots & n \end{pmatrix}$$

$$\frac{m(m+1)}{2} + \frac{m(m+1)}{2} + \dots + \frac{m(m+1)}{2} \quad \left| \quad \frac{n(n+1)}{2} + \frac{n(n+1)}{2} + \dots + \frac{n(n+1)}{2} \right.$$

$$\text{for } n \text{ columns} = \frac{n \cdot m(m+1)}{2} \quad \left| \quad \text{for } m \text{ rows} = \frac{m \cdot n(n+1)}{2} \right. \quad \text{p}$$

Solⁿ

$$\frac{mn(m+1)}{2} + \frac{mn(n+1)}{2}$$

$$\frac{mn}{2} [m+1+n+1]$$

$$\frac{mn}{2} [m+n+2]$$

Problem 8 :- If $A = (a_{ij})_{3 \times 3}$, $B = (b_{ij})_{3 \times 3}$ such that

$$b_{ij} = 2^{i+j} a_{ij} \quad \forall i, j; \quad |A| = 2; \quad |B| = ?$$

$$a) 2^{10} \quad b) 2^{11} \quad c) 2^{12} \quad \checkmark d) 2^{13}$$

$$|A| = \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} = 2$$

$$|B| = \begin{vmatrix} 2^2 a_{11} & 2^3 a_{12} & 2^4 a_{13} \\ 2^3 a_{21} & 2^4 a_{22} & 2^5 a_{23} \\ 2^4 a_{31} & 2^5 a_{32} & 2^6 a_{33} \end{vmatrix}$$

$$= 2^2 \cdot 2^3 \cdot 2^4 \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix}$$

$$= 2^9 \cdot 2 \cdot 2^2 \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix}$$

$\frac{n(n-1)}{2}$


$$= 2^{12} \cdot 2 = \underline{2^{13}}$$

Problem 4:- If $A = (a_{ij})_{n \times n}$ such that

$$i) \ a_{ij} = i^2 - j^2, \quad \forall i, j$$

$$ii) \ a_{ij} = i - j, \quad \forall i, j$$

Find sum of all elements of A



$$i) \ A = \begin{pmatrix} 0 & -3 & -8 & \dots & (1^2 - n^2) \\ 3 & 0 & -5 & \dots & (2^2 - n^2) \\ 8 & 5 & 0 & \dots & (3^2 - n^2) \\ \vdots & & & & \\ (n^2 - 1^2) & (n^2 - 2^2) & (n^2 - 3^2) & \dots & 0 \end{pmatrix}$$

A is skew symmetric matrix.

→ In skew symmetric matrix all diagonal elements must be zero & non-diagonal elements should be real no.

Benefit :- Sum of all elements of skew symmetric matrix is always zero.

Ex.

$$i) \ A = \begin{pmatrix} 0 & -3 \\ 3 & 0 \end{pmatrix}_{2 \times 2} \quad \text{Sum of all elements of skew sym. matrix} = 0$$

$$ii) \ A = \begin{pmatrix} 0 & -3 & -8 \\ 3 & 0 & -5 \\ 8 & 5 & 0 \end{pmatrix}_{3 \times 3} \quad \text{Sum} = 0$$

→ If i th row, j th column elem. of a matrix is in the form $a_{ij} = i^n - j^n$ ($n \geq 0$) then corresponding matrix is always skew symm.

* Note 3 :-

A matrix A is said to be symmetric if
 $A^T = A$ or $a_{ij} = a_{ji}$

* Note 4 :-

A matrix A is said to be skew symmetric if
 $A^T = -A$ or $a_{ij} = -a_{ji}$

UWA 2001
Problem 1 :-

The value of $\begin{vmatrix} 1+b & b & 1 \\ b & 1+b & 1 \\ 1 & 2b & 1 \end{vmatrix} = ?$

$$C_1 \rightarrow C_1 + C_2$$

$$\begin{vmatrix} 1+2b & b & 1 \\ 1+2b & 1+b & 1 \\ 1+2b & 2b & 1 \end{vmatrix} = 0$$

proportional

Problem 2 :- Find the determinant of $\begin{vmatrix} 1/a & a & bc \\ 1/b & b & ca \\ 1/c & c & ab \end{vmatrix}$

$$= \begin{vmatrix} bc/abc & a & bc \\ ca/abc & b & ca \\ ab/abc & c & ab \end{vmatrix}$$

$$= \frac{1}{abc} \begin{vmatrix} bc & a & bc \\ ca & b & ca \\ ab & c & ab \end{vmatrix} = 0$$

same

→ If there are no numbers inside determinant try to make 1 particular row or column same

classmate

Date _____

Page _____

Problem 8 :- find the value of

x	a	a	a
a	x	a	a
a	a	x	a
a	a	a	x

$$C_1 \rightarrow C_1 + C_2 + C_3 + C_4$$

$x + 3a$	a	a	a
$x + 3a$	x	a	a
$x + 3a$	a	x	a
$x + 3a$	a	a	x

$x + 3a$	1	a	a	a
	1	x	a	a
	1	a	x	a
	1	a	a	x

$$R_2 \rightarrow R_2 - R_1; \quad R_3 \rightarrow R_3 - R_1; \quad R_4 \rightarrow R_4 - R_1$$

$x + 3a$	1	a	a	a
	0	$x - a$	0	0
	0	0	$x - a$	0
	0	0	0	$x - a$

according to pt IV

$$= (x + 3a) \{ 1 \times (x - a) \times (x - a) \times (x - a) \}$$

$$= (x + 3a) (x - a)^3$$

* Short cut method :-

Procedure :-

→ If the diagonal elements are one category of same elements & non diagonal elements are other category of same elements

- 2) select the 1st row
 3) add all elements of 1st row
 4) take the product of $(1^{st} - 2^{nd}) (1^{st} - 3^{rd}) (1^{st} - 4^{th}) \dots$
 5) $(3) \times (4)$

i.e.,

 $|A| =$

$$\begin{vmatrix} x & a & a & a \\ a & x & a & a \\ a & a & x & a \\ a & a & a & x \end{vmatrix}$$

Problem 2: The value of $A = \begin{pmatrix} 1+x & 2 & 3 & 4 \\ 1 & 2+x & 3 & 4 \\ 1 & 2 & 3+x & 4 \\ 1 & 2 & 3 & 4+x \end{pmatrix}$

$$|A| = \begin{vmatrix} x+10 & 2 & 3 & 4 \\ x+10 & 2+x & 3 & 4 \\ x+10 & 2 & 3+x & 4 \\ x+10 & 2 & 3 & 4+x \end{vmatrix} = (x+10) \begin{vmatrix} 1 & 2 & 3 & 4 \\ 1 & 2+x & 3 & 4 \\ 1 & 2 & 3+x & 4 \\ 1 & 2 & 3 & 4+x \end{vmatrix}$$

$$R_2 \rightarrow R_2 - R_1 \quad R_3 \rightarrow R_3 - R_1 \quad R_4 \rightarrow R_4 - R_1$$

$$= (x+10) \begin{vmatrix} x+10 & 2 & 3 & 4 \\ 0 & x & 0 & 0 \\ 0 & 0 & x & 0 \\ 0 & 0 & 0 & x \end{vmatrix}$$

$$= (x+10) 1 \times x \times x \times x$$

$$= \underline{\underline{x^3 (x+10)}}$$

Prob

Prob

Problem 2:

$$A = \begin{vmatrix} 1+a & b & c & d \\ a & 1+b & c & d \\ a & b & 1+c & d \\ a & b & c & 1+d \end{vmatrix}$$

$$C_1 \rightarrow C_1 + C_2 + C_3 + C_4$$

$$= \begin{vmatrix} 1+a+b+c+d & b & c & d \\ 1+a+b+c+d & 1+b & c & d \\ 1+a+b+c+d & b & 1+c & d \\ 1+a+b+c+d & b & c & 1+d \end{vmatrix}$$

$$= (1+a+b+c+d) \begin{vmatrix} 1 & b & c & d \\ 1 & 1+b & c & d \\ 1 & b & 1+c & d \\ 1 & b & c & 1+d \end{vmatrix}$$

$$R_2 \rightarrow R_2 - R_1 ; R_3 \rightarrow R_3 - R_1 ; R_4 \rightarrow R_4 - R_1$$

$$= (1+a+b+c+d) \begin{vmatrix} 1 & b & c & d \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{vmatrix}$$

$$= 1+a+b+c+d \begin{bmatrix} 1 & \pm \end{bmatrix}$$

$$= 1+a+b+c+d$$

Problem 3: Find the value of

$$\begin{vmatrix} 1 & 1 & 1 \\ a & b & c \\ a^2 & b^2 & c^2 \end{vmatrix}$$

$$C_2 \rightarrow C_2 - C_1 \quad C_3 \rightarrow C_3 - C_1$$

$$A = \begin{vmatrix} \oplus & \ominus & \oplus \\ 1 & 0 & 0 \\ a & b-a & c-a \\ a^2 & b^2-a^2 & c^2-a^2 \end{vmatrix}$$

$$= 1 \left[(b-a)(c^2-a^2) - (c-a)(b^2-a^2) \right]$$

$$= (b-a)(c+a)(c-a) - (c-a)(b+a)(b-a)$$

$$= (b-a)(c-a) [c+a-b-a]$$

$$= (b-a)(c-a)(c-b)$$

$$= \{-(a-b)\} (c-a) \{-(b-c)\}$$

$$= \underline{(a-b)(b-c)(c-a)}$$

* Short cut method [for (2×2) (3×3) ... $(n \times n)$]

If matrix is in the form

$$\begin{vmatrix} 1 & 1 & 1 & 1 & \dots \\ a & b & c & d & \dots \\ a^2 & b^2 & c^2 & d^2 & \dots \\ a^3 & b^3 & c^3 & d^3 & \dots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{vmatrix}$$

then select the 2^{nd} row

& take $(2^{\text{nd}} - 1^{\text{st}})$ $(3^{\text{rd}} - 2^{\text{nd}})$... $(\text{last} - 1^{\text{st}})$

$$\text{i.e., } (a-b)(b-c)(c-d)$$

* Inverse of a matrix *

Let $A = (a_{ij})$ be $n \times n$ matrix

i) Minor: Minor of an element a_{ij} is denoted by M_{ij} and is defined as

$M_{ij} = (n-1)^{\text{th}}$ order determinant.

ii) Cofactor: Cofactor of an element a_{ij} is denoted by A_{ij} and is defined as

$$A_{ij} = (-1)^{i+j} \cdot M_{ij}$$

Ex. Consider the matrix

$$A = \begin{pmatrix} 1 & 1 & 2 \\ 2 & \textcircled{3} & 4 \\ 4 & 7 & -2 \end{pmatrix}$$

Consider the 2nd row 2nd element

$$a_{22} = 3$$

→ Minor of a_{22} is

$$M_{22} = \begin{vmatrix} 1 & 2 \\ 4 & -2 \end{vmatrix} = -2 - 8 = -10$$

→ Cofactor of a_{22} is

$$A_{22} = (-1)^{2+2} \times (-10) = -10$$

* Invertible matrix:

A matrix A is said to be invertible if we can find some other matrix B such that $AB = BA = I$ then B is called inverse of matrix A .

$$\begin{aligned}
 AB &= I \\
 A^{-1}AB &= A^{-1}I \\
 IB &= A^{-1}I \\
 \boxed{B} &= \boxed{A^{-1}}
 \end{aligned}$$

* Singular matrix:-

A matrix is said to be singular if $|A| = 0$

* Non Singular matrix:-

A matrix is said to be non singular if $|A| \neq 0$
Inverse of matrix exists if $|A| \neq 0$.

Note 1:-

$$A^{-1} = \frac{\text{adj. } A}{|A|}$$

$$\Rightarrow \text{adj. } A = |A| \cdot A^{-1}$$

$$\Rightarrow A \cdot \text{adj. } A = A \cdot |A| \cdot A^{-1}$$

$$\Rightarrow A \cdot \text{adj. } A = |A| \cdot I = \text{adj. } A \cdot A$$

Note 2:-

$$A^{-1} = \frac{\text{adj. } A}{|A|}$$

$$\Rightarrow (\text{adj. } A)^{-1} \cdot A^{-1} = (\text{adj. } A)^{-1} \cdot \frac{\text{adj. } A}{|A|}$$

$$\Rightarrow (\text{adj. } A)^{-1} \cdot A^{-1} = \frac{I}{|A|} \quad \therefore A \cdot A^{-1} = I$$

$$\Rightarrow (\text{adj. } A)^{-1} \cdot A^{-1} \cdot A = \frac{I \cdot A}{|A|}$$

$$\Rightarrow (\text{adj. } A)^{-1} = \frac{A}{|A|}$$

No

Note 3: Let A be an $n \times n$ matrix.

We know that

$$\text{adj } A = |A| \cdot A^{-1}$$

$$\Rightarrow |\text{adj } A| = \begin{matrix} |A| & A^{-1} \\ \downarrow & \downarrow \\ k & A \end{matrix}$$

$$|kA| = k^n |A|$$

$$\Rightarrow |A|^n \cdot |A^{-1}|$$

$$\Rightarrow |A|^n \cdot |A|^{-1}$$

$$\therefore |\text{adj } A| = |A|^{n-1}$$

Replacing A by $\text{adj } A$ in the above relation

$$|\text{adj } \text{adj } A| = |\text{adj } A|^{n-1} \\ = \{ |A|^{n-1} \}^{n-1}$$

$$\therefore |\text{adj } \text{adj } A| = |A|^{(n-1)^2}$$

Similarly,

$$|\text{adj } \text{adj } \text{adj } A| = |A|^{(n-1)^3}$$

& so on...

Note 4: We know that

$$A \cdot \text{adj } A = |A| \cdot I$$

replacing A by $\text{adj } A$

$$\Rightarrow \text{adj } A (\text{adj } \text{adj } A) = |\text{adj } A| \cdot I$$

$$\Rightarrow \text{adj } A (\text{adj } \text{adj } A) = |A|^{n-1} \cdot I$$

pre multiply both side by A

$$\Rightarrow A \cdot \text{adj } A (\text{adj } \text{adj } A) = A \cdot |A|^{n-1} \cdot I$$

$$|A \cdot I| = |A|$$

$$\Rightarrow |A| \cdot I (\text{adj } \text{adj } A) = |A|^{n-1} \cdot A$$

$$\Rightarrow (A) \cdot (\text{adj} \cdot \text{adj} \cdot A) = |A|^{n-1} \cdot A$$

$$\Rightarrow \text{adj} \cdot \text{adj} A = \frac{|A|^{n-1} \cdot A}{|A|}$$

$$\Rightarrow \text{adj} \cdot \text{adj} A = |A|^{n-2} A$$

Note 5:- A matrix A is said to be orthogonal if

$$A \cdot A^T = A^T \cdot A = I$$

$$A \cdot A^T = I$$

$$A^T \cdot A \cdot A^T = A^T \cdot I$$

$$I \cdot A^T = A^T \cdot I$$

$$A^T = A^{-1}$$

Note 6:- If A is an orthogonal matrix then A^{-1} & A^T are also orthogonal matrices.

Grade 2010
math 1m
problem 1

For the matrix $M = \begin{pmatrix} 3/5 & 4/5 \\ x & 3/5 \end{pmatrix}$ such that

$M^T = M^{-1}$. The value of x is ?

$$\frac{3}{5}(x) + \frac{4}{5}\left(\frac{3}{5}\right) = 0$$

$$x = -4/5$$

Note : If A is an orthogonal matrix then its rows & columns are pair wise orthogonal.
But converse of the stmt. may or may not be true.
i.e., If rows & columns of matrix are pair wise orthogonal, then the matrix may or may not be orthogonal.

Problem 2: If $A = (a_{ij})_{5 \times 5}$ such that

i) $a_{ij} = i - j, \quad \forall i, j$

ii) $a_{ij} = i^2 - j^2, \quad \forall i, j$

find A^{-1} in each case.

$$\begin{aligned} \Rightarrow i) \quad a_{ij} &= i - j \\ a_{ji} &= j - i \\ &= -(i - j) \\ &= -a_{ij} \end{aligned}$$

$$\therefore \boxed{a_{ij} = -a_{ji}} \quad \therefore A \Rightarrow \text{skew symmetric matrix}$$

$$\therefore \Rightarrow A^T = -A$$

$$\Rightarrow |A^T| = |-A|$$

$$\Rightarrow = (-1)^5 |A|$$

$$\text{as } |kA| = k^n |A|$$

$$\therefore |A^T| = -1 |A|$$

$$|A| = -|A|$$

$$|A| + |A| = 0$$

$$2|A| = 0$$

$$2 \neq 0$$

$$\therefore |A| = 0$$

$\therefore A^{-1}$ does not exist.

$|A| = |A^T| \rightarrow \text{property No 1}$

Note: ① Inverse of any odd order skew symm. matrix does not exist.

Reason: Since every odd order skew symm matrix is singular i.e., $|A| = 0$

② Inverse of even order skew symm. matrix exists.

Reason: Since every even order skew symm. matrix is non singular i.e., $|A| \neq 0$.

ii)

$$A = \begin{pmatrix} 0 & -3 \\ 3 & 0 \end{pmatrix}_{2 \times 2}$$

$$A^T = \begin{pmatrix} 0 & 3 \\ -3 & 0 \end{pmatrix} = - \begin{pmatrix} 0 & -3 \\ 3 & 0 \end{pmatrix}$$

$$A^T = -A$$

$$\therefore |A| = \begin{vmatrix} 0 & -3 \\ 3 & 0 \end{vmatrix} = 0 + 9 = (3)^2$$

⇒ The determinant of even order skew symm. matrix is a perfect square.

P. 20

⇒ If i^{th} row, j^{th} column element of a matrix is in the form

$a_{ij} = i^n - j^n$ ($n > 0$), the corresponding matrix is always skew symmetric.

Grade 10
Instu 2m
Problem 3:-

If X and Y are two non zero matrices of the same order such that $XY = (0)_{n \times n}$, then

- A) $|X| \neq 0$, $|Y| = 0$
- B) $|X| = 0$, $|Y| \neq 0$
- C) $|X| \neq 0$; $|Y| \neq 0$
- ✓ D) $|X| = 0$, $|Y| = 0$

$$XY = 0$$

take determinants on b.s

$$|XY| = 0$$

$$|X||Y| = 0$$

then $|X| = 0$ or $|Y| = 0$ or both $|X| = 0$ & $|Y| = 0$

∴ C is omitted

Let $|X| = 0$, $|Y| \neq 0 \therefore Y^{-1} \Rightarrow \text{exist}$

$$XY = 0 \Rightarrow XYY^{-1} = 0 \cdot Y^{-1}$$

$$\Rightarrow XI = 0$$

$$\Rightarrow X = 0 \quad (\text{contradiction to hypothesis})$$

$$\therefore |Y| = 0$$

If we choose $|Y| = 0$, $|X| \neq 0$ then we get

$$|X| = 0$$

$$\therefore |X| = |Y| = 0$$

Problem 4: If A, B, C, D, E, F, G are non-singular matrices of the same order such that $CEDBGAFA = I$ then B^{-1} is —



$$CEDBGAFA = I$$

$$C^{-1}CEDBGAFA = C^{-1}I$$

$$IEDBGAFA = C^{-1}I$$

$$EDBGAFA = C^{-1}$$

$$E^{-1}EDBGAFA = E^{-1}C^{-1}$$

$$IDBGAFA = E^{-1}C^{-1}$$

$$D^{-1}IDBGAFA = D^{-1}E^{-1}C^{-1}$$

$$BGAFA = D^{-1}E^{-1}C^{-1}$$

$$BGAFF^{-1} = D^{-1}E^{-1}C^{-1}F^{-1}$$

$$BGA = D^{-1}E^{-1}C^{-1}F^{-1}$$

$$BGA A^{-1} = D^{-1}E^{-1}C^{-1}F^{-1}A^{-1}$$

$$BG = D^{-1}E^{-1}C^{-1}F^{-1}A^{-1}$$

$$BG G^{-1} = D^{-1}E^{-1}C^{-1}F^{-1}A^{-1}G^{-1}$$


$$B = D^{-1}E^{-1}C^{-1}F^{-1}A^{-1}G^{-1}$$

$$\therefore B^{-1} = (D^{-1}E^{-1}C^{-1}F^{-1}A^{-1}G^{-1})^{-1}$$

$$B^{-1} = GAFCED$$

* Short cut method :-

$$CEDBGAFA$$



$$= GAFCED$$

$$\begin{array}{ccccccc} C & E & D & B & G & A & F \\ & \swarrow & & \uparrow & & \uparrow & \\ & & & A^{-1} & & & \end{array} = F C E D B G$$

$$\begin{array}{ccccccc} C & E & D & B & G & A & F \\ & & & \uparrow & & & \\ & & & F^{-1} & & & \end{array} = C E D B G A$$

Prob No. 6. Let k be a true real no. & let

$$A = \begin{pmatrix} 2k-1 & 2\sqrt{k} & 2\sqrt{k} \\ 2\sqrt{k} & 1 & -2k \\ -2\sqrt{k} & 2k & -1 \end{pmatrix}_{3 \times 3} \quad B = \begin{pmatrix} 0 & 2k-1 & \sqrt{k} \\ 1-2k & 0 & 2\sqrt{k} \\ -\sqrt{k} & -2\sqrt{k} & 0 \end{pmatrix}_{3 \times 3}$$

find i) $\det(\text{adj. } B)$ ii) $\det(\text{adj. } A)$
iii) If $\det(\text{adj. } A) = 10^6$, the value of $k = ?$

\Rightarrow i) $| \text{adj. } B | = | B |^{n-1} = 0$
 \therefore determinant of odd order skew symm. matrix is zero.

ii) $| \text{adj. } A | = | A |^{n-1} = | A |^{3-1} = | A |^2 \quad \text{--- eq}^n (1)$

$$| A | = \begin{vmatrix} 2k-1 & 2\sqrt{k} & 2\sqrt{k} \\ 2\sqrt{k} & 1 & -2k \\ -2\sqrt{k} & 2k & -1 \end{vmatrix}$$

$$R_2 \rightarrow R_2 + R_3$$

$$= \begin{vmatrix} 2k-1 & 2\sqrt{k} & 2\sqrt{k} \\ 0 & 1+2k & -2k-1 \\ -2\sqrt{k} & 2k & -1 \end{vmatrix} \quad \text{--- } \times (1+2k)$$

$$C_2 \rightarrow C_2 + C_3$$

$$= \begin{vmatrix} 2k-1 & 4\sqrt{k} & 2\sqrt{k} \\ 0 \ominus & 0 \oplus & -(1+2k) \ominus \\ -2\sqrt{k} & 2k-1 & -1 \end{vmatrix}$$

$$\Rightarrow - \left\{ - (1+2k) \left((2k-1)^2 + 8k \right) \right\}$$

$$-(1+2k) - (2k+1)(2k-1)$$

$$\Rightarrow (1+2k) (4k^2 - 4k + 1 + 8k)$$

$$\Rightarrow (1+2k) (2k^2 + 4k + 1)$$

$$\Rightarrow (2k+1) (2k+1)^2 = (2k+1)^3$$

put in eqⁿ (1)

$$\Rightarrow \left| (2k+1)^3 \right|^2 = (2k+1)^6$$

iii)

$$|\text{adj. } A| = 10^6$$

$$(2k+1)^6 = 10^6$$

$$2k+1 = 10$$

$$2k = 9$$

$$\therefore \boxed{k = 9/2}$$

Problem No. 5: If A, B, C, D are non singular matrices of the same order such that $ABCD = I$ then B^{-1} is ?

$$\overbrace{A \ B \ C \ D}^{B^{-1}} = I$$

$$\therefore \boxed{B^{-1} = CDA}$$

Prob. No. 7:—

Find the inverse of foll. matrices —

EE-05
2m 1)

$$A = \begin{pmatrix} 1 & 0 & -1 \\ 2 & 1 & -1 \\ 2 & 3 & 2 \end{pmatrix}$$

EE-95
2m H.W.
3)

$$A = \begin{pmatrix} 1 & -1 & 0 \\ 1 & 1 & -1 \\ c & 0 & 1 \end{pmatrix}$$

EE-98
2m H.W.
2)

$$A = \begin{pmatrix} 5 & 0 & 2 \\ 0 & 3 & 0 \\ 2 & 0 & 1 \end{pmatrix}$$

⇒

$$A = \begin{pmatrix} 1 & 0 & -1 \\ 2 & 1 & -1 \\ 2 & 3 & 2 \end{pmatrix}$$

$$A^{-1} = \frac{\text{adj. } A}{|A|}$$

$$\Rightarrow |A| = \begin{vmatrix} 1 & 0 & -1 \\ 2 & 1 & -1 \\ 2 & 3 & 2 \end{vmatrix}$$

$$= 1[2+3] - 1[6-2] = 5 - 4 = 1$$

$$\Rightarrow \text{adj. } A =$$

$$\begin{pmatrix} 1 & 0 & -1 \\ 2 & 1 & -1 \\ 2 & 3 & 2 \end{pmatrix} \begin{matrix} \rightarrow \text{III} \\ \rightarrow \text{I} \\ \rightarrow \text{II} \end{matrix}$$

$$\begin{pmatrix} 1 & 3 & 0 & 1 \\ -1 & 2 & -1 & -1 \\ 2 & 2 & 1 & 2 \\ 1 & 3 & 0 & 1 \end{pmatrix} \therefore \text{adj. } A = \begin{pmatrix} 5 & -3 & 1 \\ -6 & 4 & -1 \\ 4 & -3 & 1 \end{pmatrix}$$

$$\therefore A^{-1} = \begin{pmatrix} 5 & -3 & 1 \\ -6 & 4 & -1 \\ 4 & -3 & 1 \end{pmatrix}$$

Procedure: (3x3)

- ⇒ Select the middle row middle element & move in anticlockwise direction to complete 1 cycle. The corresponding element will be return in the 1st column separately.

2) Select the 3rd row middle element & move in anti-clockwise direction to complete 1 cycle.
The corresponding element will be return in the 2nd column separately.

3) Repeat the same process with 1st row also.

4) Copy 1st column as a last column & find det. of smaller matrices.

Procedure: (2x2)

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

$$A^{-1} = \frac{1}{ad-bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$$

EC-2008
2m

Prob. No. 8:— Given an orthogonal matrix

$$A = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & 0 & 0 \\ 0 & 0 & 1 & -1 \end{pmatrix} \text{ then } (A A^T)^{-1} \text{ is } = ?$$

$$\text{Orthogonal} \Rightarrow (A A^T) = I$$

$$(A A^T)^{-1} = I^{-1}$$

but Inverse / adj. of Identity matrix is Identity matrix only.

$$\therefore (A A^T)^{-1} = I$$

Prob. No. 9 Find the inverse of $A = \begin{pmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{pmatrix}$

→ Here A is an orthogonal matrix therefore.

$$A^{-1} = A^T$$

* Rank of Matrix *

* Sub matrix : A matrix obtained by deleting some rows or columns or both is called as sub matrix.

Ex. Consider the matrix

$$A = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 1 & -1 & 0 \end{pmatrix}$$

$$B = \begin{pmatrix} 1 & 2 \\ 4 & 5 \end{pmatrix} \quad B_2 = \begin{pmatrix} 2 & 3 \\ 5 & 6 \end{pmatrix} \quad B_3 = \begin{pmatrix} 4 & 5 & 6 \\ 1 & -1 & 0 \end{pmatrix}$$

sub matrices of A

* Minor : The determinant of square sub matrix is called its minor.

Ex.

$$\left. \begin{aligned} |B_1| &= \begin{vmatrix} 1 & 2 \\ 4 & 5 \end{vmatrix} = 5 - 8 = -3 \\ |B_2| &= \begin{vmatrix} 2 & 3 \\ 5 & 6 \end{vmatrix} = 12 - 15 = -3 \end{aligned} \right\} \text{Minors of A}$$

* Rank : If the determinant of highest possible square matrix is not equal to zero then the order of the determinant is called Rank of matrix.

Ex. Find the rank of

$$A = \begin{pmatrix} 1 & 3 & -2 & 1 \\ 2 & -1 & 4 & 2 \\ 1 & -11 & 14 & 5 \end{pmatrix}_{3 \times 4}$$

$$\begin{vmatrix} 1 & 3 & -2 \\ 2 & -1 & 4 \\ 1 & -11 & 14 \end{vmatrix}_{3 \times 3} = 1[-14 + 44] - 3[28 - 4] - 2[-22 + 1] \\ = 30 - 72 + 42 = 0$$

$$\begin{vmatrix} 3 & -2 & 1 \\ -1 & 4 & 2 \\ -11 & 14 & 5 \end{vmatrix}_{3 \times 3} = 3[20 - 28] + 2[-5 + 22] + 1[-14 + 44] \\ = -24 + 34 + 30 = 40 \neq 0$$

\therefore Rank of $A = \rho(A) = 3$ i.e., order of matrix

* Row Echelon Form:-

A matrix A is said to be in Row Echelon form iff

- i) zero rows should occupy the last rows, if any.
- ii) the no. of zero's before a non zero element of each row is less than no. of such zeros before a non zero element of the next row.

Ex.

$$A = \begin{pmatrix} 0 & 1 & 2 & 3 & 4 & 0 \\ 0 & 0 & -1 & 0 & 2 & 0 \\ 0 & 0 & 0 & 4 & 5 & 0 \\ 0 & 0 & 0 & 7 & 9 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}_{6 \times 6}$$

Condition 1 satisfy
——— 2 ———

Note: The rank of Row Echelon form matrix is equal to no. of non zero rows.

$$\boxed{\rho(A) = 4} \leftarrow \text{Linearly Independent Row/Vector}$$

- These non zero rows are called Linearly Independent rows/vector.
- To reduce any matrix into row echelon form we should use only row operations.
- Every upper triangular matrix is in R.E. form but every R.E. form will not be an upper triangular matrix.

Mo

Ex. 1

$$A = \begin{pmatrix} 1 & 2 & 3 \\ 0 & -4 & 2 \\ 0 & 0 & 7 \end{pmatrix}$$

↓
Upper Δ^{lan} matrix
 $\therefore A$ is in R.E. form

Ex. 2

$$A = \begin{pmatrix} 0 & 2 & 3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

↓
Not Upper Δ^{lan} matrix
but A is in R.E. form

Upper Δ^{lan} matrix \Rightarrow R.E. form R.E. form \nRightarrow Upper Δ^{lan} matrix

* Column Echelon form:—

A matrix A is said to be in column echelon form iff

i) zero ~~rows~~ columns should occupy the last columns, if any.

ii) The no. of zeros above a non zero element of each column is less than the no. of zeros above a

non zero element of the next column.

Ex. Consider the matrix

$$A = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \\ 1 & 2 & 0 & 0 & 0 \\ 1 & 2 & 3 & 0 & 0 \\ 1 & 2 & 3 & 4 & 0 \end{pmatrix}$$

Note :-

Rank of matrix in Column Echelon form is equal to no. of non zero columns.

$$\therefore \boxed{R(A) = 4}$$

→ To reduce any matrix into column echelon form, we should use only Column Operations

→ Every lower Δ^{th} matrix will be in column echelon form.

but every C.E. form will not be a lower Δ^{th} matrix.

Ex. 1

$$A = \begin{pmatrix} 1 & 0 & 0 \\ 3 & 5 & 0 \\ 9 & 4 & 5 \end{pmatrix}$$



Lower Δ^{th} matrix

Lower Δ^{th} M \Rightarrow C.E. form

Ex. 2

$$A = \begin{pmatrix} 0 & 0 & 0 \\ 1 & 1 & 1 \\ 2 & 0 & 0 \end{pmatrix}$$

C.E. form \nRightarrow Lower Δ^{th} matrix

Example: Consider the matrix $A = \begin{pmatrix} 1 & 3 & -2 \\ 2 & -1 & 4 \\ 1 & -11 & 14 \end{pmatrix}$

\Rightarrow Row Echelon form

$$R_2 \rightarrow R_2 - 2R_1 \quad R_3 \rightarrow R_3 - R_1$$

$$A = \begin{pmatrix} 1 & 3 & -2 \\ 0 & -7 & 8 \\ 0 & -14 & 16 \end{pmatrix} \quad R_3 \rightarrow R_3 - 2R_2$$

$$A = \begin{pmatrix} 1 & 3 & -2 \\ 0 & 7 & 8 \\ 0 & 0 & 0 \end{pmatrix} \quad \therefore \boxed{\rho(A) = 2}$$

\Rightarrow Column Echelon form

$$C_2 \rightarrow C_2 - 3C_1 \quad C_3 \rightarrow C_3 + 2C_1$$

$$A = \begin{pmatrix} 1 & 0 & 0 \\ 2 & -7 & 8 \\ 1 & -14 & 16 \end{pmatrix} \quad C_3 \rightarrow 7C_3 + 8C_2$$

$$A = \begin{pmatrix} 1 & 0 & 0 \\ 2 & -7 & 0 \\ 1 & -14 & 0 \end{pmatrix} \quad \therefore \boxed{\rho(A) = 2}$$

Note: Rank of matrix = no. of non zero rows &
no. of non zero columns.

Prob. No. 7

2)

$$A = \begin{pmatrix} 5 & 0 & 2 \\ 0 & 3 & 0 \\ 2 & 0 & 1 \end{pmatrix}$$

$$A^{-1} = \frac{\text{adj. } A}{|A|}$$

$$|A| = \begin{vmatrix} 5 & 0 & 2 \\ 0^{\ominus} & 3^{\oplus} & 0^{\ominus} \\ 2 & 0 & 1 \end{vmatrix} = 3[5-4] = 3$$

$$\text{adj. } |A| = \begin{pmatrix} 3 & 0 & 0 & 3 \\ 0 & 1 & 2 & 0 \\ 0 & 2 & 5 & 0 \\ 3 & 0 & 0 & 3 \end{pmatrix}$$

$$\text{adj. } A = \begin{pmatrix} 3 & 0 & -6 \\ 0 & 1 & 0 \\ -6 & 0 & 15 \end{pmatrix}$$

$$\therefore A^{-1} = \frac{1}{3} \begin{pmatrix} 3 & 0 & -6 \\ 0 & 1 & 0 \\ -6 & 0 & 15 \end{pmatrix}$$

$$3) \quad A = \begin{pmatrix} 1 & -1 & 0 \\ 1 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$$

$$A^{-1} = \frac{\text{Adj. } A}{|A|}$$

$$|A| = \begin{vmatrix} 1 & -1 & 0 \\ 1 & 1 & 1 \\ 0^{\oplus} & 0^{\ominus} & 1^{\oplus} \end{vmatrix} = 1[1+1] = 2$$

$$\text{adj. } A = \begin{pmatrix} 1 & 0 & -1 & 1 \\ 1 & 1 & 0 & 1 \\ 1 & 0 & 1 & 1 \\ 1 & 0 & -1 & 1 \end{pmatrix}$$

$$= \begin{pmatrix} 1 & 1 & -1 \\ -1 & 1 & -1 \\ 0 & 0 & 2 \end{pmatrix}$$

$$\therefore A^{-1} = \frac{1}{2} \begin{pmatrix} 1 & 1 & -1 \\ -1 & 1 & -1 \\ 0 & 0 & 2 \end{pmatrix}$$

* Information regarding Rank of matrix :

- 1) $\rho(O_{n \times n}) = 0$
- 2) $\rho(I_{n \times n}) = n$
- 3) $\rho\{\text{adj. } I_{n \times n}\} = n$
- 4) $\rho(A) = \rho(A^T)$
- 5) $\rho(A+B) \leq \rho(A) + \rho(B)$
- 6) $\rho(A-B) \geq \rho(A) - \rho(B)$
- 7) $\rho(AB) \geq \rho(A) + \rho(B) - n$, if A & B are $n \times n$ matrix
- 8) $\rho(AB) \leq \min\{\rho(A), \rho(B)\}$
- 9) If A is an $m \times n$ matrix, then $\rho(A) \leq \min(m, n)$
- 10) If $\rho(A_{n \times n}) = 0$, then $\rho(\text{adj. } A) = 0$
- 11) If $\rho(A_{n \times n}) = n-1$, then $\rho(\text{adj. } A) = 1$
- 12) If $\rho(A_{n \times n}) = n-2$, then $\rho(\text{adj. } A) = 0$

^{2m}
Problem 1: If $A = (a_{ij})_{m \times n}$, such that $a_{ij} = i \cdot j, \forall i, j$
 then $\rho(A) = ?$

$$A = \begin{pmatrix} 1 & 2 & 3 & \dots & n \\ 2 & 4 & 6 & \dots & 2n \\ 3 & 6 & 9 & \dots & 3n \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ m & 2m & 3m & \dots & mn \end{pmatrix}_{m \times n}$$

to find $\rho(A)$, convert the matrix into Row Echelon form

$$R_2 \rightarrow R_2 - 2R_1, \quad R_3 \rightarrow R_3 - 3R_1, \quad R_m \rightarrow R_m - mR_1$$

$$A = \begin{pmatrix} 1 & 2 & 3 & \dots & n \\ 0 & 0 & 0 & \dots & 0 \\ 0 & 0 & 0 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & 0 \end{pmatrix}$$

$\therefore \rho(A) = \text{No. of non zero rows} = 1$

Problem 2: If $x = (x_1, x_2, \dots, x_n)^T$ is n -tuple non zero vector then

i) $\rho(x x^T)$

ii) $\rho(x^T x)$

i) $\rho(x x^T)$

$$x = (x_1, x_2, \dots, x_n)^T$$

$$x = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}_{n \times 1} \quad x^T = (x_1, x_2, \dots, x_n)_{1 \times n}$$

$$\rho(x x^T) \leq \min \{ \rho(x), \rho(x^T) \}$$

$$\rho(x x^T) \leq \min \{ 1, 1 \}$$

$$\Rightarrow \rho(x x^T) \leq 1 \quad \begin{matrix} \swarrow \\ 0 \end{matrix} \quad \begin{matrix} \searrow \\ x \end{matrix} \quad (x \rightarrow \text{Non zero vector})$$

$$\therefore \boxed{\rho(x x^T) = 1}$$

mate

Problem 3: The rank of 5×6 matrix Q is 4 then which of the foll. stmt is true.

- a) Q will have 4 L.I rows & 4 L.I columns
- b) Q will have 4 L.I rows & 5 L.I columns
- c) $Q Q^T$ is invertible
- d) $Q^T Q$ is invertible

$$\rho(Q_{5 \times 6}) = 4$$

\therefore 4 non zero rows or columns

option (a)

option (b) : X

If det of matrix $\neq 0$ then order of matrix

classmate

Date

Page

option (c) $(Q)_{5 \times 6} \cdot (Q)^T_{6 \times 5}$

$$(QQ^T)_{5 \times 5} \rightarrow \text{Invertible}$$

$$(QQ^T)^{-1} \text{ exists}$$

$$|(QQ^T)_{5 \times 5}| \neq 0 \Rightarrow \rho(QQ^T) = 5 \text{ (contradiction to stmt)}$$

option (d) $(Q^T)_{6 \times 5} \cdot (Q)_{5 \times 6}$

$$(Q^T Q)_{6 \times 6} \rightarrow \text{Invertible}$$

$$(Q^T Q)^{-1} \text{ exists}$$

$$|(Q^T Q)_{6 \times 6}| \neq 0 \Rightarrow \rho(Q^T Q) = 6 \text{ (contradiction to stmt)}$$

* Linearly Dependent & Independent vectors *

⇒ Two vectors x_1 & x_2 are L.D. if one vector is expressed as multiple of other vector.

x_1 & $x_2 \Rightarrow$ same directional vectors


Example

$$x_1 = \begin{pmatrix} 1 \\ 2 \end{pmatrix}, \quad x_2 = \begin{pmatrix} 3 \\ 6 \end{pmatrix}$$

$$x_2 = 3 \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

$$\boxed{x_2 = 3x_1} \quad \text{or} \quad \boxed{x_1 = \frac{1}{3}x_2}$$

$$\therefore x_1, x_2 \Rightarrow \text{L.D.}$$



$\vec{1} \quad \vec{2} \quad \vec{3} \quad \vec{4} \quad \vec{5} \rightarrow x_1$

⇒ Two vectors in R^2 are L.D. if and only if they are collinear.

⇒ Three vectors in R^3 are L.D. iff they are coplanar.

⇒ Two vectors x_1, x_2 are L.I. iff it is not possible to express one vector as a multiple of other vector.

Example:

$$x_1 = \begin{pmatrix} 1 \\ 2 \end{pmatrix} \quad x_2 = \begin{pmatrix} 3 \\ 4 \end{pmatrix}$$

$$x_1 \neq k x_2 \quad \text{OR} \quad x_2 \neq k x_1$$

$$\therefore x_1, x_2 \Rightarrow \text{L.I.}$$

* Linearly Dependent vectors: (for 2 or more than 2 vectors)

A set of r n -vectors x_1, x_2, \dots, x_r are said to be linearly dependent if there exist r scalars k_1, k_2, \dots, k_r such that

$$k_1 x_1 + k_2 x_2 + \dots + k_r x_r = 0$$

where k_1, k_2, \dots, k_r not all zeros.

(at least one k value is a non zero no.)

* Linearly Independent vectors:

A set of r n -vectors x_1, x_2, \dots, x_r are said to be linearly independent if there exist r scalar k_1, k_2, \dots, k_r such that

$$k_1 x_1 + k_2 x_2 + \dots + k_r x_r = 0$$

all zeros.

* Criteria for L.I & L.D

- 1) If $\rho(A) = \text{no. of given vectors}$ or $|A| \neq 0$, the given vectors are said to be L.I.

Ex.

Consider the vectors

$$x_1 = (1 \ 2 \ 2), \quad x_2 = (2 \ 1 \ 2), \quad x_3 = (2 \ 2 \ 1)$$

$$|A| = \begin{vmatrix} 1 & 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 & 1 \end{vmatrix} = 1(1-4) - 2(2-4) + 2(4-2) \\ = -3 + 4 + 4 \\ = 5$$

$$\therefore |A| = 5 \neq 0 \Rightarrow \text{L.I.}$$

- 2) If $\rho(A) < \text{no. of given vectors}$ or $|A| = 0$, the given vectors are said to be L.D.

Ex.

Consider the vectors

$$x_1 = (1 \ 3 \ -2), \quad x_2 = (2 \ -1 \ 4), \quad x_3 = (1 \ -11 \ 14)$$

$$|A| = \begin{vmatrix} 1 & 3 & -2 \\ 2 & -1 & 4 \\ 1 & -11 & 14 \end{vmatrix} = 1(-14+44) - 3(28-4) - 2(-22+1) \\ = 30 - 72 + 42$$

$$|A| = 0 \Rightarrow \text{L.D.}$$

- 3) If the given vectors are L.D. then any one of the vector can be expressed as linear combination of other vectors.

- 4) If the given vectors are L.I then it is not possible to write any one of the vector as linear combination of other vectors.

at least one elem. must be non zero

5) Every non zero vector is L.I. vector.

Ex.

Consider the vector

$$x = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$$

$$kx = 0$$

$$k \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

Non
zero
vector

$$\therefore k = 0$$

$\therefore x \rightarrow$ L.I. vector

columns/rows of Identity matrix

6) The set of unit vectors are always L.I.

Ex. Consider the set of vectors

$$x_1 = (1, 0, 0), x_2 = (0, 1, 0), x_3 = (0, 0, 1)$$

$$|A| = \begin{vmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{vmatrix} = 1$$

$$|A| = 1 \neq 0 \Rightarrow \text{L.I. vectors}$$

7) The set of vectors having at least one zero vector are L.D.

Ex. Consider the set of vectors

$$x_1 = (1, 2, 3), x_2 = (0, 0, 0), x_3 = (1, -1, 4)$$

$$|A| = \begin{vmatrix} 1 & 2 & 3 \\ 0 & 0 & 0 \\ 1 & -1 & 4 \end{vmatrix} = 0 \quad \therefore x_1 = x_2 = x_3 \Rightarrow \text{L.D.}$$

$$|A| = 0 \Rightarrow \text{L.D.}$$

No. of vectors = r

No. of elements in each vector = n

classmate

Date

Page

- 8) A set of r vectors with $r \leq n$ components (elements) are always L.I., provided the vectors should not be in the same direction.

Ex. Consider the vectors

$$(1 \ -1 \ 2 \ 1), (1 \ 2 \ 3 \ 4), (2 \ 3 \ 4 \ 9)$$

$$r = 3$$

$$n = 4$$

here, $r < n \Rightarrow$ L.I.

- 9) A set of r vectors with $r > n$ components then given vectors are L.D.

Ex. consider the vectors

$$x_1 = (1 \ 1 \ 2)$$

$$x_2 = (1 \ -1 \ 0)$$

$$x_3 = (1 \ -1 \ 5)$$

$$x_4 = (9 \ -5 \ 4)$$

$$r = 4$$

$$n = 3$$

here, $r > n \Rightarrow$ L.D.

$\therefore x_1, x_2, x_3, x_4$ are L.D.

- 10) A set of r vectors with $r = n$ components may be L.I or L.D.

* Dimension & Basis of the vectors

* Dimension :- It is defined as no. of L.I. vectors.

Dimension = No. of L.I. vectors = no. of non zero rows in Row Echelon form = no. of non zero columns in Column Echelon form.

* Basis :- It is defined as the set of L.I. vectors.

Basis = set of L.I. vectors = set of non zero rows in R.E. form = set of non zero columns in C.E. form.

Problem 1: Test whether the following vectors are L.D or L.I.
Also find their dimension & basis.

$$\begin{pmatrix} 1 & 1 & -1 & 0 \end{pmatrix}, \begin{pmatrix} 4 & 4 & -3 & 1 \end{pmatrix}, \begin{pmatrix} -6 & 2 & 2 & 2 \end{pmatrix}, \begin{pmatrix} 9 & 9 & -6 & 3 \end{pmatrix}$$

$m = 4$ $n = 4$ here $r = n$ \therefore may be L.D or L.I

$$A = \begin{vmatrix} 1 & 1 & -1 & 0 \\ 4 & 4 & -3 & 1 \\ -6 & 2 & 2 & 2 \\ 9 & 9 & -6 & 3 \end{vmatrix}$$

$$R_2 \rightarrow R_2 - 4R_1 \quad R_3 \rightarrow R_3 + 6R_1 \quad R_4 \rightarrow R_4 - 9R_1$$

$$A = \begin{vmatrix} 1 & 1 & -1 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 8 & -4 & 2 \\ 0 & 0 & 3 & 3 \end{vmatrix}$$

$$-3+4$$

$$2+6 \quad 2+6$$

$$-6+9$$

$$R_2 \leftrightarrow R_3$$

$$A = \begin{vmatrix} 1 & 1 & -1 & 0 \\ 0 & 8 & -4 & 2 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 3 & 3 \end{vmatrix}$$

$$R_4 \rightarrow R_4 - 3R_3$$

$$A = \begin{vmatrix} 1 & 1 & -1 & 0 \\ 0 & 8 & -4 & 2 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 \end{vmatrix} \quad \left. \begin{array}{l} \\ \\ \\ \end{array} \right\} 3 \text{ L.I. Vectors}$$

→ whenever a matrix is reduced to RREF form then Rank of matrix = Dimension of Set of L.I. vectors are called basis.

CLASSMATE
Date: / /
Page: /

$$\rho(A) = 3 < \text{No. of given vectors (4)}$$

$$\therefore x_1, x_2, x_3, x_4 \Rightarrow \text{L.I.}$$

$$\left[\begin{array}{cccc|c} 1 & 1 & -1 & 0 & 0 \\ 0 & 8 & -4 & 2 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right] \Rightarrow \text{rank} \neq 0$$

$$\text{Dimension} = 3$$

$$\text{Basis} = \{ (1 \ 1 \ -1 \ 0), (0 \ 8 \ -4 \ 2), (0 \ 0 \ 1 \ 1) \}$$

Gate (2m)

Problem: If q_1, q_2, \dots, q_m are n -dimensional vectors with $m < n$. The vectors are L.I. The matrix Q is q_1, q_2, \dots, q_m as columns. The rank of $Q = ?$

a) m b) n c) between m & n ✓ m

$$q_1 = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}$$

$$Q = \begin{pmatrix} | & | & | & \dots & | \\ q_1 & q_2 & q_3 & & q_m \\ | & | & | & & | \end{pmatrix}_{n \times m} \Rightarrow (Q)_{n \times m}$$

$$\rho(Q_{n \times m}) \leq \min(n, m)$$

but $m < n$ — given

$$\therefore \rho(Q_{n \times m}) \leq m$$

$$\therefore \boxed{\rho(Q) < m}$$

$m \times$ from criteria (2)
 $< m$ ✓

* Nullity of a matrix :-

Page

* Nullity :- It is denoted by $N(A)$.

& is defined as the difference betⁿ order of matrix and rank of matrix.

i.e.,

$$\boxed{N(A) = n(A) - \rho(A)}$$

\downarrow \downarrow \downarrow
 Nullity order Rank

\Rightarrow Nullity of a non-singular matrix is always zero.

Let A be an $n \times n$ non singular matrix.

Then,

$$|A_{n \times n}| \neq 0$$

$$\therefore \rho(A) = n$$

$$N(A) = n(A) - \rho(A)$$

$$= n - n$$

$$\therefore N(A) = 0$$

oblem: The nullity of $A = \begin{pmatrix} k & 1 & 2 \\ 1 & -1 & -2 \\ 1 & 1 & 4 \end{pmatrix} = 4$. The value of $k = ?$

$$N(A) = n(A) - \rho(A)$$

$$1 = 3 - \rho(A)$$

$$\rho(A) = 2$$

for 3×3 matrix

iff $\rho(A) = 3$ then $|A| \neq 0$

$$|A| = 0 \Rightarrow$$

$$\begin{vmatrix} k & 1 & 2 \\ 1 & -1 & -2 \\ 1 & 1 & 4 \end{vmatrix} = 0$$

$$k = -1$$

$$ax + by + cz = 2 \implies \text{Non-Homogeneous}$$

Date _____

Page

$$N(A) = n(A) - p(A)$$

$$3 = n(A) - 5$$

$$n(A) = 8$$

Consider the following non homogeneous system of m linear eq^{ns} in n unknowns.

$$\left. \begin{aligned} a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n &= b_1 \\ a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n &= b_2 \\ \vdots &\vdots \\ a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n &= b_m \end{aligned} \right\} \text{I}$$

Procedure

→ Write the given system of eq^{ns} in the form $AX = B$

i.e.,

$$\begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} = \begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{pmatrix}$$

$\begin{matrix} \uparrow & & \uparrow & & \uparrow \\ \text{coefficient matrix} & \text{Sol}^n \text{ matrix} & \text{column matrix} \\ & & & & \text{of constants.} \end{matrix}$

2) Write the elements of matrix B in the last column of matrix A. The resulting matrix is called Augmented matrix & is denoted by $(A|B)$

$$(A|B) = \left(\begin{array}{cccc|c} a_{11} & a_{12} & \dots & a_{1n} & b_1 \\ a_{21} & a_{22} & \dots & a_{2n} & b_2 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} & b_m \end{array} \right)$$

- 3) Reduce the augmented matrix $(A|B)$ into Row Echelon form & hence find rank of A & rank of $(A|B)$
- 4) If $\rho(A) < \rho(A|B)$ or $\rho(A|B) \neq \rho(A)$, the given system of equations ~~are said to~~ have no solⁿ (inconsistent). No
- 5) If $\rho(A|B) = \rho(A) = \text{no. of unknowns}$, the given system of eqⁿ have unique solution.
- 6) If $\rho(A|B) = \rho(A) < \text{no. of unknowns}$, the given system of eqⁿ have infinite no. of solutions.
- 7) If the given system of equations have a solⁿ (unique or infinite solⁿ).
The solⁿ can be found by reducing the matrix AB into Row Echelon form & by using back code substitution, the variables x_1, x_2, \dots, x_n can be found. Proof

Note: If the total no. of eq^{ns} $<$ total no. of variables, the given system of eqⁿ have infinite no. of sol^{ns}. Proof

These infinite no. of sol^{ns} can be found by assigning $(n-r)$ variables as arbitrary constants.
These $(n-r)$ sol^{ns} are linearly independent sol^{ns}.

$$x + y = 3$$

$$r = 1, \quad n = 2$$

$$r < n, \quad n - r = 2 - 1 = 1$$

put $y = c \rightarrow$ L.I solⁿ

$$x + y = 3$$

$$x + c = 3$$

$$x = 3 - c$$

$$x = 3 - c$$

$$y = c$$

solⁿ Note: Consider the system of eq^{ns}

$$ax + by = e$$

$$cx + dy = f$$

The above system of eq^{ns} have

1) No solⁿ if $\frac{a}{c} = \frac{b}{d} \neq \frac{e}{f}$

2)

2) Unique solⁿ if $\frac{a}{c} \neq \frac{b}{d}$

3) Infinite solⁿ if $\frac{a}{c} = \frac{b}{d} = \frac{e}{f}$

ECE2010
(1M)

Problem 1: The system of eq^{ns}

$$4x + 2y = 7$$

$$2x + y = 6$$

have

$$\frac{4}{2} = \frac{2}{1} \neq \frac{7}{6}$$

$$\left(\frac{a}{c} = \frac{b}{d} \neq \frac{e}{f} \right)$$

\therefore The given system of eq^{ns} have no solⁿ

Problem 2:

$$x + 2y = 5$$

$$2x + 3y = 9$$

$$\frac{1}{2} \neq \frac{2}{3} \neq \frac{5}{9}, \quad \frac{1}{2} \neq \frac{2}{3} \left(\frac{a}{c} \neq \frac{b}{d} \right)$$

\therefore Unique solⁿ

Problem B: $x + y = 3$

$$3x + 5y = 9$$

$$\frac{1}{3} \times 3 = \frac{3}{9} \therefore \text{Infinite sol}^n$$

Problem 4: How many solutions does the following system of eq^{ns} have

$$-x + 5y = -1$$

$$x - y = 2$$

$$x + 3y = 3$$

a) Infinite b) exactly 2 c) Unique solⁿ d) No solⁿ

$$(A|B) = \left(\begin{array}{cc|c} \textcircled{-1} & 5 & -1 \\ 1 & -1 & 2 \\ 1 & 3 & 3 \end{array} \right)$$

$$R_2 \rightarrow R_2 + R_1$$

$$R_3 \rightarrow R_3 + R_1$$

$$= \left(\begin{array}{cc|c} -1 & 5 & -1 \\ 0 & \textcircled{4} & 1 \\ 0 & 8 & 2 \end{array} \right)$$

$$R_3 \rightarrow R_3 - 2R_2$$

$$= \left(\begin{array}{cc|c} -1 & 5 & -1 \\ 0 & 4 & 1 \\ 0 & 0 & 0 \end{array} \right)$$

A B

$$\rho(A|B) = 2$$

$$\rho(A) = 2$$

$$\rho(A|B) = \rho(A) = \text{No. of unknowns} = 2$$

\therefore Unique solⁿ

(2m)

Problem 5: Consider following non homogeneous system of Linear eqⁿ in 3 variables x_1, x_2, x_3 .

$$2x_1 - x_2 + 3x_3 = 1$$

$$3x_1 + 2x_2 + 5x_3 = 2$$

$$-x_1 + 4x_2 + x_3 = 3$$

The above system of eqⁿ have —

- a) No solⁿ b) ☒ unique solⁿ c) more than 1 but finite no of sol^{ns} d) Infinite no of sol^{ns}

$$(A|B) = \begin{pmatrix} 2 & -1 & 3 & 1 \\ 3 & 2 & 5 & 2 \\ -1 & 4 & 1 & 3 \end{pmatrix}$$

$$R_2 \rightarrow 2R_2 - 3R_1$$

$$R_3 \rightarrow 2R_3 + R_1$$

$$(A|B) = \begin{pmatrix} 2 & -1 & 3 & 1 \\ 0 & 7 & 1 & 1 \\ 0 & 7 & 5 & 7 \end{pmatrix}$$

$$R_3 \rightarrow R_3 - R_2$$

$$(A|B) = \begin{pmatrix} 2 & -1 & 3 & 1 \\ 0 & 7 & 1 & 1 \\ 0 & 0 & 4 & 6 \end{pmatrix}$$

$$\rho(A|B) = 3$$

$$\rho(A) = 3$$

No. of unknowns = 3

∴ Unique solⁿ

2010 (2m)

Problem 6: For the set of eq^{ns} $x_1 + 2x_2 + x_3 + x_4 = 2$

$$3x_1 + 6x_2 + 3x_3 + 3x_4 = 6$$

which of the foll. stmt is true —

- 1) There exist only trivial solⁿ
 2) There are no sol^{ns}

3) Unique non trivial solⁿ non zero solⁿ

4) Infinite no of non trivial solⁿ

\Rightarrow No. of eq^{ns} (m) = 2
 No. of variables (n) = 4
 here, $m < n$
 Infinite solⁿ (Non trivial)

Problem 7: The value of x_3 obtained by solving the foll. system of eq^{ns}.

$$x_1 + 2x_2 - 2x_3 = 4$$

$$2x_1 + x_2 + x_3 = -2$$

$$-x_1 + x_2 - x_3 = 2$$

$$(A|B) = \left(\begin{array}{ccc|c} 1 & 2 & -2 & 4 \\ 2 & 1 & 1 & -2 \\ -1 & 1 & -1 & 2 \end{array} \right)$$

$$R_2 \rightarrow R_2 - 2R_1$$

$$R_3 \rightarrow R_3 + R_1$$

$$(A|B) = \left(\begin{array}{ccc|c} 1 & 2 & -2 & 4 \\ 0 & -3 & 5 & -10 \\ 0 & 3 & -3 & 6 \end{array} \right)$$

$$R_3 \rightarrow R_3 + R_2$$

$$(A|B) = \left(\begin{array}{ccc|c} 1 & 2 & -2 & 4 \\ 0 & -3 & 5 & -10 \\ 0 & 0 & 2 & -4 \end{array} \right)$$

$$2x_3 = -4$$

$$\boxed{x_3 = -2}$$

(To find x_2

$$-3x_2 + 5x_3 = -10$$

$$-3x_2 - 10 = -10$$

$$-3x_2 = 0$$

$$\boxed{x_2 = 0}$$

2011
(2m)

Problem 8: find the values of λ, u for which the following system of eq^{ns}

$$x + y + z = 6$$

$$x + 4y + 6z = 20$$

$$x + 4y + \lambda z = u \quad \text{have}$$

1) No solⁿ

2) infinite solⁿ

3) Unique solⁿ

$$(A|B) = \left(\begin{array}{ccc|c} \textcircled{1} & 1 & 1 & 6 \\ & 1 & 4 & 20 \\ & 1 & 4 & \lambda & u \end{array} \right)$$

$$R_2 \rightarrow R_2 - R_1$$

$$R_3 \rightarrow R_3 - R_1$$

$$(A|B) = \left(\begin{array}{ccc|c} & 1 & 1 & 6 \\ & 0 & \textcircled{2} & 14 \\ & 0 & 3 & \lambda - 1 & u - 6 \end{array} \right)$$

$$R_3 \rightarrow R_3 - R_2$$

$$(A|B) = \left(\begin{array}{ccc|c} & 1 & 1 & 6 \\ & 0 & 3 & 14 \\ & 0 & 0 & \lambda - 6 & u - 20 \end{array} \right)$$

$$\rho(A) = 3$$

$$\rho(A|B) = 3$$

1) No solⁿ : If $\rho(A) < \rho(A|B)$

If $\lambda = 6, u \neq 20$

$$\rho(A) = 2$$

$$\rho(A|B) = 3$$

$\Rightarrow \rho(A) < \rho(A|B) \rightarrow \text{No solⁿ}$

2) Unique solⁿ : If $\rho(A) = \rho(A|B) = \text{No. of unknowns}$

For $\lambda \neq 6$ & $u \rightarrow \text{any value}$ if $u = 20$ or $u \neq 20$

3) Infinite no. of solⁿ :- If $r(A) = r(A|B) < \text{No. of unknowns}$

No. of unknowns = 3

must be 2

$r(A)$ & $r(A|B)$ should be less than 3 bcz rank can't be 0 or 1

It is possible if

$$r = 0 \quad \& \quad u = 20 \implies \text{Infinite sol}^n \text{ set}$$

HLW

Problem 9: Find the values of A & B for which the following system of eq^{ns}

$$x + 2y + 3z = 6$$

$$x + 3y + 5z = 9$$

$$2x + 5y + \alpha z = \beta \quad \text{have}$$

1) No solⁿ 2) Unique solⁿ 3) Infinite sol^{ns}

Grade
20

Problem 10: For what values of α & β , the following system of eq^{ns}

$$x + y + z = 5$$

have infinite no

$$x + 3y + 3z = 9$$

of solⁿ?

$$\alpha + 2y + \alpha z = \beta$$

a) $\alpha = 2, \beta = 7$

c) $\alpha = 3, \beta = 4$

b) $\alpha = 7, \beta = 2$

d) $\alpha = 4, \beta = 3$

$$(A|B) = \begin{pmatrix} \textcircled{1} & 1 & 1 & 5 \\ 1 & 3 & 3 & 9 \\ 1 & 2 & \alpha & \beta \end{pmatrix}$$

$$R_2 \rightarrow R_2 - R_1$$

$$R_3 \rightarrow R_3 - R_1$$

$$= \begin{pmatrix} 1 & 1 & 1 & 5 \\ 0 & \textcircled{2} & 2 & 4 \\ 0 & 1 & \alpha-1 & \beta-5 \end{pmatrix}$$

$$R_3 \rightarrow 2R_3 - R_2$$

Non-homogeneous system eqⁿ \Rightarrow draw augmented matrix

classmate

Date _____

Page _____

$$A = \begin{pmatrix} 1 & 1 & 1 & 5 \\ 0 & 2 & 2 & 4 \\ 0 & 0 & 2\alpha - 4 & 2\beta - 14 \end{pmatrix}$$

$$\rho(A) = \rho(A|B) < \text{No. of unknowns}$$

$$\rho(A) = \rho(A|B) < 3$$

$$2\alpha - 4 = 0$$

$$2\beta - 14 = 0$$

$$2\alpha = 4$$

$$2\beta = 14$$

$$\boxed{\alpha = 2}$$

$$\boxed{\beta = 7}$$

Problem II : If A is 3×4 matrix & the non homogeneous system of equations $AX=B$ is inconsistent (No sol). The highest possible rank of A is

\Rightarrow

$$(A|B)_{3 \times 5}$$

$$(A|B)_{3 \times 5}$$

$$\rho\{(A|B)\}_{3 \times 5} \leq \min(3, 5)$$

$$\rho\{(A|B)\} \leq 3$$

$= 3$ Highest Possible Rank

But it is given that the given system of eq^{ns} are inconsistent (No solⁿ)

$$\text{for inconsistent} \rightarrow \rho(A) < \rho(A|B)$$

$$\rho(A) < 3$$

Highest possible rank of $A = \underline{\underline{2}}$

$$\rho(A) \begin{cases} 2 \\ 1 \\ 0 \end{cases}$$

* Homogeneous system of Linear Eqⁿ *

Consider the following homogeneous system of Linear eq^s in m equations & n unknowns

$$\left. \begin{aligned} a_{11}x_1 + a_{12}x_2 + a_{13}x_3 + \dots + a_{1n}x_n &= 0 \\ a_{21}x_1 + a_{22}x_2 + a_{23}x_3 + \dots + a_{2n}x_n &= 0 \\ \vdots &\vdots \\ a_{m1}x_1 + a_{m2}x_2 + a_{m3}x_3 + \dots + a_{mn}x_n &= 0 \end{aligned} \right\} \textcircled{I}$$

Procedure to solve problems :-

1) Write the given system of eq^s in the form $AX=0$

$$\begin{pmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \dots & a_{2n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & a_{m3} & \dots & a_{mn} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{pmatrix}$$

2) Reduce the matrix A into either row or column echelon form (but always row echelon form is preferable) OR find the determinant of matrix $|A|$.

3) If $\rho(A) = \text{No. of unknowns (variables)}$ OR $|A| \neq 0$ ($A \rightarrow$ Non singular matrix), the given system of eq^s have trivial solⁿ.
($x=y=z=0$)

4) If $\rho(A) < \text{No. of unknowns}$ OR $|A| = 0$, the given system of eq^s have infinite no. of solⁿ.


no. of solⁿ are dependent on $n-r$

classmate

Date

Page

All these infinite no. of solⁿ can be found by assigning $(n-r)$ variables as arbitrary constant. These $(n-r)$ solⁿs are called L.I. solⁿs



Problem I: for what values of λ the system of eq^s

$$x + y + z = 0$$

$$(\lambda + 1)x + y + (\lambda + 1)z = 0$$

$$(\lambda^2 - 1)z = 0$$

have 2 L.I. solⁿ ?



$$\text{No. of L.I. solⁿ} = n - r = 2$$

$$= 3 - r = 2$$

$$\therefore \boxed{r = 1} \rightarrow \text{Rank of matrix}$$

For $\lambda = 1$

Let $x = 1, z = 1$

$$|A| = \begin{vmatrix} 1 & 1 & 1 \\ 2+1 & \lambda+1 & \lambda+1 \\ 0 & 0 & \lambda^2-1 \end{vmatrix} = 0 \Rightarrow \text{Upper } \Delta^{\text{an}}$$

Prob1

$$\Rightarrow (\lambda+1)(\lambda^2-1) = 0$$

$$(\lambda+1)(\lambda+1)(\lambda-1) = 0$$

$$\lambda = -1, 1 \dots \text{not correct}$$

put in matrix in order to get

$$|A| = 1$$

$$\therefore \boxed{\lambda = -1}$$

Prob

Problem 2:- The rank of 3×3 matrix A is 1. The homogeneous system of eq^{ns} $AX=0$ has

- a) trivial solⁿ b) 1 L.T. solⁿ
 c) 2 L.T. sol^{ns} d) 3 L.T. sol^{ns}

$$(A)_{3 \times 3} \Rightarrow 3 \text{ eq}^{\text{ns}} \text{ \& } 3 \text{ variables}$$

$$n = 3$$

$$r = 1 \Rightarrow r < n$$

$$\Downarrow$$
Infinite solⁿ

$$\therefore \text{L.T. sol}^{\text{ns}} = n - r$$

$$= 3 - 1$$

$$= 2$$

2011 (pm)

Problem 3:- The system of eq^{ns}: $2x_1 + x_2 + x_3 = 0$

$$x_2 - x_3 = 0$$

$$x_1 + x_2 = 0 \quad \text{have}$$

- a) No non trivial solⁿ c) 5 Non trivial sol^{ns}
 b) Unique non trivial solⁿ d) Infinite non trivial sol^{ns}

$$\begin{vmatrix} 2 & 1 & 1 \\ 0 & 1 & -1 \\ 1 & 1 & 0 \end{vmatrix} = 2[1] - 1[0+1] + 1[0-1]$$

$$= 2 - 1 - 1 = 0$$

Infinite solⁿ

Problem 4: The system of eqⁿ

$$\begin{aligned} x + 3y - 2z &= 0 \\ 2x - y + 4z &= 0 \\ x - 11y + 14z &= 0 \end{aligned}$$

$$\begin{vmatrix} 1 & 3 & -2 \\ 2 & -1 & 4 \\ 1 & -11 & 14 \end{vmatrix} = \begin{aligned} &+[-14+44] - 2[28-4] - 1[-22+1] \\ &= 30 - 48 + 42 \end{aligned}$$

$$1[-14+44] - 3[28-4] + 2[-22+1]$$

$$= 0 \quad = 30 - 72 + 42 = 0$$

Problem 5: Find the values of k for which the following system of eq^{ns} have infinite no. of non trivial sol^{ns}.

$$(3k-8)x + 3y + 3z = 0$$

$$3x + (3k-8)y + 3z = 0$$

$$3x + 3y + (3k-8)z = 0$$

$$|A| = 0$$

→ If $|A| = 0$, the given system of eq^{ns} have infinite no. of non zero sol^{ns}.

$$\begin{vmatrix} 3k-8 & 3 & 3 \\ 3 & 3k-8 & 3 \\ 3 & 3 & 3k-8 \end{vmatrix} = 0$$

$$\Rightarrow (3k-8+3+3)(3k-8-3)(3k-8-3) = 0$$

$$(3k-2)(3k-11)(3k-11) = 0$$

$$3k = 2$$

$$3k = 11$$

$$\therefore k = \frac{2}{3}, \frac{11}{3}$$

Problem 6 Find the real value of λ for which the following system of eq^{ns} have non trivial sol^{ns}

infinite no. of non trivial solⁿ

$$x + 2y + 3z = \lambda x$$

$$3x + y + 2z = \lambda y$$

$$2x + 3y + z = \lambda z$$

$$\lambda = 6$$

Imp

* Eigen values & Eigen Vectors *

$$A = \begin{pmatrix} 4 & 2 \\ 2 & 4 \end{pmatrix}$$

$$A - \lambda I = \begin{pmatrix} 4 & 2 \\ 2 & 4 \end{pmatrix} - \lambda \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

$$= \begin{pmatrix} 4 & 2 \\ 2 & 4 \end{pmatrix} - \begin{pmatrix} \lambda & 0 \\ 0 & \lambda \end{pmatrix}$$

\mathbb{R}

$$A - \lambda I = \begin{pmatrix} 4-\lambda & 2 \\ 2 & 4-\lambda \end{pmatrix}$$

↓

characteristic matrix

ch. det

ch. polynomial

ch. eqⁿ

$$|A - \lambda I| = \begin{vmatrix} 4-\lambda & 2 \\ 2 & 4-\lambda \end{vmatrix} = (4-\lambda)^2 - 4$$

$$= \lambda^2 - 8\lambda + 16 - 4$$

$$|A - \lambda I| = 0 \implies \lambda^2 - 8\lambda + 12 = 0$$

$$\therefore \lambda = 6, 2$$

↓

ch. roots or
Eigen values

Null vector can not be Eigen vector

CLASSMATE

Date

Page

* Eigen values: Let A be an $n \times n$ matrix. λ is a scalar (some constant). The matrix $A - \lambda I$ is called as characteristic matrix.

$|A - \lambda I|$ is called characteristic determinant or characteristic polynomial.

The roots of this ch. det. are called char roots or Eigen values or Latent roots or Proper values.

The set of eigen values of matrix A is called as spectrum of A .

* Eigen Vector: If λ is an eigen value of a matrix A then there exist a non zero vector X such that $AX = \lambda X$, then the non zero vector X is called as Eigen vector.

Note:

$$AX = \lambda X$$

$$AX = \lambda XI$$

$$AX - \lambda XI = 0$$

$$AX - \lambda IX = 0$$

$$(A - \lambda I) X = 0$$

Trivial solⁿ $\Rightarrow |A - \lambda I| \neq 0$

Infinite solⁿ $\Rightarrow |A - \lambda I| = 0$

Non-trivial solⁿ

\Downarrow
Eigen Vectors

Note: Consider the matrix

$$A = \begin{pmatrix} 4 & 2 \\ 2 & 4 \end{pmatrix}$$


$$\begin{pmatrix} 4 & 2 \\ 2 & 4 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 4+2 \\ 2+4 \end{pmatrix} = \begin{pmatrix} 6 \\ 6 \end{pmatrix} = 6 \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

\downarrow \downarrow \uparrow \uparrow
 A X λ X
 (non zero vector)

$X = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$ is eigen vector corresponding to eigen value $\lambda = 6$ for matrix $A = \begin{pmatrix} 4 & 2 \\ 2 & 4 \end{pmatrix}$

→ Any pt./vector exist along the same vector will also be an eigen vector corresponding to the same eigen value.

Ex. (2,2) (3,3) ... (7,7) (8,8) ...



Eq $A = \begin{pmatrix} 4 & 2 \\ 2 & 4 \end{pmatrix}$

$$\begin{pmatrix} 4 & 2 \\ 2 & 4 \end{pmatrix} \begin{pmatrix} 1 \\ 2 \end{pmatrix} = \begin{pmatrix} 4+4 \\ 2+8 \end{pmatrix} = \begin{pmatrix} 8 \\ 10 \end{pmatrix} = 2 \begin{pmatrix} 4 \\ 5 \end{pmatrix}$$

\downarrow \downarrow \downarrow
 A X X

Not same

$x = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$ is not eigen vector corresponding to eigen value 2

Problem: Find the eigen values & eigen vector of

$$A = \begin{pmatrix} 3 & 4 \\ 4 & -3 \end{pmatrix}$$

Symmetric matrix

$$|A - \lambda I| = 0$$

$$\begin{vmatrix} 3-\lambda & 4 \\ 4 & -3-\lambda \end{vmatrix} = (3-\lambda)(-3-\lambda) - 16$$

$$= -9 - 3\lambda + 3\lambda + \lambda^2 - 16 = 0$$

$$\lambda^2 - 25 = 0$$

$$\lambda^2 = 25$$

$$\lambda = \pm 5$$

\therefore Eigen values = 5, -5

case i)

$$(A - \lambda I) X = 0$$

$$\begin{pmatrix} 3-\lambda & 4 \\ 4 & -3-\lambda \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

↑ Eigen vectors

put $\lambda = 5$

$$\begin{pmatrix} -2 & 4 \\ 4 & -8 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$-2x_1 + 4x_2 = 0 \quad \text{--- (1)}$$

$$4x_1 - 8x_2 = 0 \quad \text{--- (2)}$$

eqⁿ (1) \Rightarrow divide b.s. by (-2)

$$x_1 - 2x_2 = 0$$

$$\frac{x_1}{x_2} = \frac{2}{1}$$

$$x_1 = 2x_2$$

$$\therefore X_1 = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 2 \\ 1 \end{pmatrix}$$

Case ii)

when $\lambda = -5$

$$(A - \lambda I) X = 0$$

$$\begin{pmatrix} 4 & 4 \\ 4 & -3-\lambda \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

put $\lambda = -5$

$$\begin{pmatrix} 8 & 4 \\ 4 & 2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$8x_1 + 4x_2 = 0 \quad \text{--- (1)}$$

$$4x_1 + 2x_2 = 0 \quad \text{--- (2)}$$

2 vectors are
not den.only 1 vector is
den

$$2x_1 + x_2 = 0$$

$$2x_1 = -x_2$$

$$\frac{x_1}{x_2} = \frac{-1}{2}$$

$$x_2 = \left(\frac{x_1}{x_2} \right) = \begin{pmatrix} -1 \\ 2 \end{pmatrix}$$

$$x_1 = \begin{pmatrix} 2 \\ 1 \end{pmatrix} \quad x_2 = \begin{pmatrix} -1 \\ 2 \end{pmatrix}$$

$$x_1^T x_2 = 0 \quad \text{or} \quad x_1 x_2^T = 0 \Rightarrow x_1 \& x_2 \text{ are orthogonal}$$

$$x_1^T \cdot x_2 = \begin{pmatrix} 2 & 1 \end{pmatrix} \begin{pmatrix} -1 \\ 2 \end{pmatrix} = 2x_1 + 1x_2$$

$$(x_1^T \cdot x_2)_{1 \times 1} = -2 + 2 = 0$$

$\therefore x_1 \& x_2 \Rightarrow$ orthogonal vectors

E.V. of Upper Δ , Lower Δ , diag... only
diagonal elem. only

classmate

Date _____
Page _____

Note 1: The Eigen vectors corresponding to diff. eigen value of a real symmetric matrix are always orthogonal.

Note 2: If the Eigen vectors corresponding to diff. eigen values of any square matrix are always linearly Independent.

$$X_1 = \begin{pmatrix} 2 \\ 1 \end{pmatrix} \quad X_2 = \begin{pmatrix} -1 \\ 2 \end{pmatrix}$$

$$X_1 \neq k X_2 \quad \text{OR} \quad X_2 \neq k X_1$$

$$\therefore \begin{matrix} X_1 \\ X_2 \end{matrix} > \text{L.I.} \quad (\text{acc}^n \text{ to criteria (4)})$$

is In the above ex. the eigen values of matrix are 5, -5 the corresponding eigen vectors are $\begin{pmatrix} 2 \\ 1 \end{pmatrix}$ & $\begin{pmatrix} -1 \\ 2 \end{pmatrix}$

* Consider the matrix $A = \begin{pmatrix} 2 & 3 \\ 0 & 2 \end{pmatrix}$

lower L.I.

$$\lambda = 2, 2 \rightarrow \text{Repeated twice}$$

$$\text{when } \lambda = 2 \quad \text{Max. no. of L.I. vectors}$$

$$(A - \lambda I)X = 0$$

$$\begin{pmatrix} 2-\lambda & 3 \\ 0 & 2-\lambda \end{pmatrix} \begin{pmatrix} X_1 \\ X_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\text{put } \lambda = 2$$

$$\begin{pmatrix} 0 & 3 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} X_1 \\ X_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$0 X_1 + 3 X_2 = 0$$

$$3 X_2 = 0$$

$$\therefore X_2 = 0$$

$$x_1 \neq 0$$

$$\text{put } x_1 = c$$

$$x = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} c \\ 0 \end{pmatrix} \text{ where } c \neq 0$$

Acc" to criteria (5)


every non zero vector is L.I vector.

Note:- If some of the eigen value of a matrix are repeated then the eigen vectors corresponding to repeated eigen values may be L.I or L.D.

If an eigen value λ is repeated n times the eigen vectors corresponding to repeated eigen values are always L.I. which are given by

$$p = n - r \quad ; \quad 1 \leq p \leq n \quad \left\{ \begin{array}{l} \text{max. } n \text{ times} \\ \text{no. of unknowns or variables} \end{array} \right\}$$

\downarrow
 $S(A - \lambda I)$



$$A_{3 \times 3}$$

$$\lambda_1, \lambda_2, \lambda_3$$

$$\lambda_1 \neq \lambda_2 \neq \lambda_3$$

$$\downarrow \quad \downarrow \quad \downarrow$$

$$x_1 \quad x_2 \quad x_3$$

L.I

$$\lambda_1 \neq \lambda_2 \neq \lambda_3$$

$$\downarrow \quad \downarrow \quad \downarrow$$

$$x_1 \quad x_2 \quad x_3$$

L.I L.I or L.D L.I

$$\lambda_1 = \lambda_2 = \lambda_3$$

$$\downarrow \quad \downarrow \quad \downarrow$$

$$x_1 \quad x_1, x_2 \quad x_1, x_2, x_3$$

$$\downarrow \quad \downarrow \quad \downarrow$$

L.I L.I L.I

$$\begin{pmatrix} c_1 + c_2 \\ c_2 \\ c_1 \end{pmatrix} = c_1 \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} + c_2 \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}$$

$$\downarrow \quad \downarrow$$

$$x_1 \quad x_2$$

L.I

2011 Grade
Instr.
(m)
Note:

$$A \cdot X = \lambda \cdot X$$

Eigen vector of A

Eigen value of A

$$A^{-1} \cdot A \cdot X = A^{-1} \cdot \lambda \cdot X$$

$$I \cdot X = A^{-1} \cdot \lambda \cdot X$$

$$X = \lambda \cdot A^{-1} X$$

$$\frac{X}{\lambda} = A^{-1} X$$

$$A^{-1} \cdot X = \frac{1}{\lambda} \cdot X$$

Eigen vector of A^{-1}

Eigen value of A^{-1}

If λ is eigen value of A & X is eigen vector of A but $\frac{1}{\lambda}$ is eigen value of A^{-1} and

X is eigen vector of A^T . Therefore, A & A^T have same eigen vectors.

A & A^m have same eigen vectors ($m > 0$) corresponding to eigen values.

$$AX = \lambda X$$

$$AAX = A\lambda X$$

$$A^2 X = \lambda \lambda X$$

$$A^2 X = \lambda^2 X$$

17/07/2012
* Properties of Eigen values & Eigen vectors:-

1) Sum of eigen values is equal to trace of matrix
Sum of diagonal elem

i.e., if $\lambda_1, \lambda_2, \dots, \lambda_n$ are eigen values of a matrix A then

① $\text{Trace } A = \lambda_1 + \lambda_2 + \dots + \lambda_n$

2) ② $|A| = \lambda_1 \cdot \lambda_2 \cdot \lambda_3 \cdot \dots \cdot \lambda_n$

Ex. Consider the matrix

$$A = \begin{pmatrix} 5 & 4 \\ 1 & 2 \end{pmatrix}$$

$$5 - \lambda \quad 4$$

$$1 \quad 2 - \lambda$$

$$(5 - \lambda)(2 - \lambda) - 4$$

$$10 - 5\lambda - 2\lambda + \lambda^2 - 4 = \lambda^2 - 7\lambda + 6$$

$$\lambda = 1, 6$$

i) $\rightarrow 1 + 6 = 5 + 2$

$\therefore \text{Trace of } A = 7$

ii) $\rightarrow 1 \times 6 = 10 - 4 = 6$

$$|A| = 6$$

3) The eigen values of upper triangular or lower triangular or diagonal or scalar or identity matrix is its diagonal elements.

$$1) A = \begin{pmatrix} 1 & 2 & 3 \\ 0 & -4 & 2 \\ 0 & 0 & 7 \end{pmatrix}; \lambda = 1, -4, 7$$

Upper Δ lar

$$2) A = \begin{pmatrix} 3 & 0 & 0 \\ 4 & 5 & 0 \\ 0 & 8 & 9 \end{pmatrix}; \lambda = 3, 5, 9$$

Lower Δ lar

$$3) A = \begin{pmatrix} 3 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 5 \end{pmatrix}; \lambda = 3, 4, 5$$

Diagonal matrix

$$4) A = \begin{pmatrix} 3 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 3 \end{pmatrix}; \lambda = 3, 3, 3$$

scalar matrix

$$5) A = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}; \lambda = 0, 0, 0$$

scalar / Null matrix

$$6) A = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}; \lambda = 1, 1, 1$$

Identity matrix

4) The eigen values of A & A^T are same

5) The eigen values of A & $P^{-1}AP$ are same where P is a non singular matrix.

6) The eigen values of real symmetric matrix are real.

7) The eigen values of skew symmetric matrix are purely imaginary or zeros.

8) The eigen values of orthogonal matrix are of unit modulus i.e., ± 1 .

9) If λ is an eigen value of an orthogonal matrix then $\frac{1}{\lambda}$ is also one of its eigen value.

10) If λ is an eigen value of matrix A then

i) $k\lambda$ is an eigen value of kA .

ii) $\frac{1}{\lambda}$ is also an eigen value of A^{-1} .

iii) λ^2 is an eigen value of A^2 .

iv) λ^m is an eigen value of A^m .

v) $\frac{|A|}{\lambda}$ is an eigen value of $\text{adj. } A$.

vi) $\lambda \pm k$ is an eigen value of $A \pm kI$.

vii) $(\lambda \pm k)^2$ is an eigen value of $(A \pm kI)^2$.

viii) $\frac{1}{\lambda \pm k}$ is an eigen value of $(A \pm kI)^{-1}$.

Note: If A is a singular matrix i.e., $|A|=0$ then one of its eigen value should be zero

$$\lambda_1 \cdot \lambda_2 \cdot \lambda_3 = |A|$$

↓

$$0 \cdot \lambda_2 \cdot \lambda_3 = |A|$$

$$|A|=0$$

$\therefore |A| \rightarrow$ singular

Problem 1: $A = \begin{pmatrix} -1 & 3 & 5 \\ -3 & -1 & 6 \\ 0 & 0 & 3 \end{pmatrix}$ are

$$j = \sqrt{-1}$$

A) $3, 3+5j, 6-j$

B) $-6+5j, 3-j, 3+j$

C) $3-j, 3+j, 5+j$

☒ D) $3, -1+3j, -1-3j$

by Prop. 1 :

$$\text{Trace } A = \lambda_1 + \lambda_2 + \lambda_3$$

[cross check]

$$-1 - 1 + 3 = 3 - 1 + 3j - 1 - 3j$$

$$1 = 1$$

If 1 or more optⁿ satisfy prop. 1

Problem 2: The eigen values

& eigen vector of a 2×2 matrix are given by

Eigen value

Eigen vector

$$\lambda_1 = 8$$

$$V_1 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

$$\lambda_2 = 4$$

$$V_2 = \begin{pmatrix} 1 \\ -1 \end{pmatrix}$$

the matrix is —

☒ a) $\begin{pmatrix} 6 & 2 \\ 2 & 6 \end{pmatrix}$

b) $\begin{pmatrix} 4 & 6 \\ 6 & 4 \end{pmatrix}$

c) $\begin{pmatrix} 2 & 4 \\ 4 & 2 \end{pmatrix}$

d) $\begin{pmatrix} 4 & 8 \\ 8 & 4 \end{pmatrix}$

to find eigen value system

$$(A - \lambda I) X = 0$$

classmate

Date _____

Page _____

1998
(2m)

Problem 3: The vector $\begin{pmatrix} 1 \\ 2 \\ -1 \end{pmatrix}$ is an eigen vector of the

P₂

matrix

$$A = \begin{pmatrix} -2 & 2 & -3 \\ 2 & 1 & -6 \\ -1 & -2 & 0 \end{pmatrix}$$

The eigen value corresponding to the eigen vector is -

$$(A - \lambda I) X = 0$$

$$\begin{pmatrix} -2-\lambda & 2 & -3 \\ 2 & 1-\lambda & -6 \\ -1 & -2 & 0-\lambda \end{pmatrix} \begin{pmatrix} 1 \\ 2 \\ -1 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

P₂

value consider any 1 row & multiplied

$$-1 - 4 + \lambda = 0$$

$$\boxed{\lambda = 5}$$

P₂

Problem 4: For the matrix $\begin{pmatrix} 4 & 2 \\ 2 & 4 \end{pmatrix}$ the eigen value corresponding to the eigen vector $\begin{pmatrix} 1 & 0 & 1 \\ 1 & 0 & 1 \end{pmatrix}$ is

$$(A - \lambda I) X = 0$$

P₂

$$\begin{pmatrix} 4-\lambda & 2 \\ 2 & 4-\lambda \end{pmatrix} \begin{pmatrix} 1 & 0 & 1 \\ 1 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

P₂

$$2 + 4 - \lambda = 0$$

$$\lambda = 6$$

Problem 5: The min & max eigen values of a matrix $\begin{pmatrix} 1 & 1 & 3 \\ 1 & 3 & 1 \\ 3 & 1 & 1 \end{pmatrix}$ are -2 & 6 resp. what would be the 3rd eigen value.

$$\begin{aligned} \text{Trace } A &= \lambda_1 + \lambda_2 + \lambda_3 \\ &= 1 + 5 + 1 = -2 + 6 + \lambda_3 \\ \boxed{\lambda_3 &= 3} \end{aligned}$$

Problem 6: The matrix $\begin{pmatrix} 1 & 2 & 4 \\ 3 & 0 & 6 \\ 1 & 1 & p \end{pmatrix}$ has an eigen value = 3. Sum of other two eigen values is —

$$\begin{aligned} 1 + 0 + p &= 3 + \lambda_2 + \lambda_3 \\ p + 1 - 3 &= \lambda_2 + \lambda_3 \\ \boxed{\lambda_2 + \lambda_3 &= p - 2} \end{aligned}$$

2011
Problem 7: Consider the following matrix

$A = \begin{pmatrix} 2 & 3 \\ x & y \end{pmatrix}$ The eigen values of A are 4 & 8. The values of x, y are

$$\begin{aligned} \text{prop. 1} &\rightarrow 2 + y = 4 + 8 \\ y &= 12 - 2 = 10 \end{aligned}$$

$$\text{prop. 2} \rightarrow 2 \times y = 20 \quad 4 \times 8 = 32$$

$$\begin{vmatrix} 2 & 3 \\ x & 10 \end{vmatrix} = 32$$

$$x = -4$$

$$20 - 3x = 32$$

$$20 - 32 = 3x$$

$$-12 = 3x \quad \boxed{x = -4}$$

$$x = -\frac{10}{3}$$

$$2y - 3x = 4 \times 8 \Rightarrow 2 \times 10 - 3x = 32$$

$$-3x = 32 - 20$$

$$-3x = 12$$

$$\boxed{x = -4}$$

Prob 8:- The eigen values of 3×3 matrix are given by 1, -3, 9. Find

i) Trace $(A^2 + A^{-1} - \text{adj. } A)$

ii) det $(A^2 + A^{-1} - \text{adj. } A)$

$$\Rightarrow |A| = 1 \times -3 \times 9 = -27$$

$$A^2 + A^{-1} - \text{adj. } A$$

$$\rightarrow \lambda^2 + \frac{1}{\lambda} - \frac{|A|}{\lambda} \quad \text{by prop. (10)}$$

$$\begin{array}{l} 1 \\ -3 \\ 9 \end{array}$$

$$\lambda^2 + \frac{1}{\lambda} - \frac{|A|}{\lambda} \quad \begin{array}{l} (1)^2 + \frac{1}{1} - \frac{(-27)}{1} = 29 \\ (-3)^2 + \frac{1}{(-3)} - \frac{(-27)}{(-3)} \\ 9 - \frac{1}{3} - 9 = -\frac{1}{3} \\ 9^2 + \frac{1}{9} - \frac{(-27)^3}{9} \end{array}$$

$$= 29 - \frac{1}{3} + 27$$

$$= 56 - \frac{1}{3}$$

$$= \frac{168 - 1}{3} = \frac{167}{3}$$

$$= \frac{167}{3}$$

$$\frac{729 + 1 + 27}{9} = \frac{757}{9}$$

$$\therefore \text{Trace } (A^2 + A^{-1} - \text{adj. } A) = 29 - \frac{1}{3} + \frac{757}{9}$$

$$ii) \det(A^2 + A^T - \text{adj } A) = 29 \times \left(\frac{-1}{3}\right) \times \frac{757}{9}$$

Prob. 9:- Given $A = \begin{pmatrix} 1 & 2 & -3 \\ 0 & 3 & 2 \\ 0 & 0 & -2 \end{pmatrix}$, the eigen values of $3A^3 + 5A^2 - 6A + 2I$ are

Upper
Triangular

$$\therefore \lambda = 1, 3, -2$$

$$3A^3 + 5A^2 - 6A + 2I \Rightarrow 3\lambda^3 + 5\lambda^2 - 6\lambda + 2$$

put $\lambda = 1$

$$3(1)^3 + 5(1)^2 - 6(1) + 2$$

$$3 + 5 - 6 + 2 = 4$$

put $\lambda = 3$

$$3 \times 27 + 5 \times 9 - 18 + 2$$

$$81 + 45 - 16 = 126 - 16 = 110$$

put $\lambda = -2$

$$-(3 \times 8) + 20 + 12 + 2$$

$$-24 + 24 = 0$$

$$3 \times (-8) + 5 \times 4 + 12 + 2$$

$$-24 + 20 + 12 + 2 = 10$$

2009
(cm)

Prob. No. 10:- The eigen values of 2×2 matrix A are given by -2 & -3 resp. the eigen values of $(x+I)^{-1} \cdot (x+5I)$

This is not possible

$$\frac{-577}{9} \Rightarrow (x+I)^{-1} (x+5I) = (x+I)^{-1} \{ (x+I) + 4I \}$$

$$= (x+I)^{-1} \cdot (x+I) + 4I \cdot (x+I)^{-1}$$

$$= I + 4(x+I)^{-1}$$

$$4(x+I)^{-1} + I \Rightarrow \frac{4}{1+\lambda} + 1$$

$$\Rightarrow \frac{4}{1+\lambda} + 1$$

$$\text{put } \lambda = -2$$

$$\frac{4}{1-2} + 1 = -3$$

$$\text{put } \lambda = -3$$

$$\frac{4}{1-3} + 1 = -1$$

20

Pro

Prob 11 :- The eigen vector of a 3×3 matrix are
are orthogonal. That will be
the 3rd orthogonal eigen vector.

$$\begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}$$

$$x_1 = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} \quad x_2 = \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix} \quad \text{Let } x_3 = \begin{pmatrix} x \\ y \\ z \end{pmatrix}$$

here, 3 vectors are orthogonal

$$x_1^T \cdot x_3 = 0 \quad x_2^T \cdot x_3 = 0$$

$$\begin{pmatrix} 1 & 0 & 1 \end{pmatrix} \cdot \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \quad \begin{pmatrix} 1 & 0 & -1 \end{pmatrix} \cdot \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

$$x + z = 0 \quad \text{--- (i)}$$

$$x - z = 0 \quad \text{--- (ii)}$$

add (i) & (ii)

$$2x = 0$$

$$\therefore x = 0 \quad \text{put in (i)}$$

$$0 + z = 0$$

$$z = 0$$

$y \neq 0$ \therefore zero vector can't be eigen vector

$$\Rightarrow y = c \quad c - \text{arbitrary constant \& } c \neq 0$$

\therefore The 3rd eigen vector is $\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$

$$x = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$$

2008
(2m)

Prob. 12 :- The eigen vector of 2×2 matrix $\begin{pmatrix} 1 & 2 \\ 0 & 2 \end{pmatrix}$ are given by $\begin{pmatrix} 1 \\ a \end{pmatrix}, \begin{pmatrix} 1 \\ b \end{pmatrix}$, what is $(a+b)$?

$\lambda = 1, 2$

$AX = \lambda X$ OR $(A - \lambda I)X = 0$

put $\lambda = 1$

$$\begin{pmatrix} 1 & 2 \\ 0 & 2 \end{pmatrix} \begin{pmatrix} 1 \\ a \end{pmatrix} = \begin{pmatrix} 1 \\ 2 \end{pmatrix} \Rightarrow \begin{pmatrix} 1 & 2 \\ 0 & 2 \end{pmatrix} \begin{pmatrix} 1 \\ a \end{pmatrix} = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

$2 \times 2 \quad 2 \times 1$

~~$1+2a = 2a$~~

$1+2a = 1$

$\& \quad 2a = 0$

$\boxed{a = 0}$

put $\lambda = 2$

$$\begin{pmatrix} 1 & 2 \\ 0 & 2 \end{pmatrix} \begin{pmatrix} 1 \\ b \end{pmatrix} = 2 \begin{pmatrix} 1 \\ b \end{pmatrix} = \begin{pmatrix} 2 \\ 2b \end{pmatrix}$$

$1+2b = 2$

$2b = 1$

$\boxed{b = 1/2}$

for

$c \neq 0$

Method I: —

Case 1: when $\lambda = 1$

$$(A - \lambda I) X = 0$$

$$\begin{pmatrix} 1-\lambda & 2 \\ 0 & 2-\lambda \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

Prob.

$$\begin{pmatrix} 0 & 2 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$2x_2 = 0 \quad \therefore x_2 = 0$$

here $x_1 \neq 0$

$x_1 = c$ $c \rightarrow$ non zero arbitrary const

$$X = \begin{pmatrix} c \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad \text{generally } c \text{ is replaced by } 1$$

Case 2: when $\lambda = 2$

$$\begin{pmatrix} 1-2 & 2 \\ 0 & 2-\lambda \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\begin{pmatrix} -1 & 2 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$-x_1 + 2x_2 = 0$$

$$2x_2 = x_1$$

$$\frac{2}{1} = \frac{x_1}{x_2}$$

$$x_2 = \begin{pmatrix} 1 \\ 1/2 \end{pmatrix}$$

$$x_1 = 2 \quad x_2 = 1$$

$$\frac{x_1}{x_2} = \frac{1}{(1/2)}$$

$$x_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ a \end{pmatrix} \Rightarrow a = 0$$

$$x_2 = \begin{pmatrix} 1 \\ 1/2 \end{pmatrix} = \begin{pmatrix} 1 \\ b \end{pmatrix} \quad b = 1/2$$

Prob. 13: find eigen values & eigen vectors of foll.

a) $A = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 2 & 2 \\ 0 & 0 & 3 \end{pmatrix}$

$$\Rightarrow \lambda = 1, 2, 3$$

→ put $\lambda = 1$

$$\begin{array}{c|ccc} x_1 & x_2 & x_3 & \\ \hline 1 & 0 & 0 & 1 \\ 1 & 2 & 0 & 1 \end{array}$$

$$\frac{x_1}{2} = \frac{x_2}{0} = \frac{x_3}{0}$$

$$x_1 = \begin{pmatrix} 2 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$$

other form of this eigenvector

→ put $\lambda = 2$

$$\begin{array}{c|cccc} x_1 & x_2 & x_3 & & \\ \hline 1 & 0 & -1 & 0 & \\ 0 & 2 & 0 & 0 & \end{array}$$

$$\frac{x_1}{2} = \frac{x_2}{2} = \frac{x_3}{0}$$

$$x_2 = \begin{pmatrix} 2 \\ 2 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}$$

for distinct eigen values

→ consider all 3 rows?

→ select any one row and solve

→ if row is all zero then

consider another row

if all rows are zero then

select any one column

→ put $x_1 = 1$ or $x_2 = 1$

→ put $\lambda = 3$

$$\begin{array}{c|cccc} x_1 & x_2 & x_3 & & \\ \hline 1 & 0 & -2 & 1 & \\ -1 & 2 & 0 & -1 & \end{array}$$

$$\frac{x_1}{2} = \frac{x_2}{4} = \frac{x_3}{2}$$

$$x_3 = \begin{pmatrix} 2 \\ 4 \\ 2 \end{pmatrix} = \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix}$$

b) $A = \begin{pmatrix} 1 & 2 & 3 \\ 0 & -4 & 2 \\ 0 & 0 & 7 \end{pmatrix}$ No

put $\lambda = 1$

$$\begin{array}{ccc} x_1 & x_2 & x_3 \\ 2 & 3 & 2 \\ 5 & 2 & 0 \end{array}$$

$$\frac{x_1}{19} = \frac{x_2}{0} = \frac{x_3}{0} \quad X_1 = \begin{pmatrix} 19 \\ 0 \\ 0 \end{pmatrix}$$

put $\lambda = -4$

$$\begin{array}{ccc} x_1 & x_2 & x_3 \\ 2 & 3 & 5 \\ -8 & 2 & 0 \end{array}$$

$$\frac{x_1}{20} = \frac{x_2}{-10} = \frac{x_3}{-40} \quad X_2 = \begin{pmatrix} 28 \\ -10 \\ -40 \end{pmatrix}$$

put $\lambda = 7$

$$\begin{array}{ccc} x_1 & x_2 & x_3 \\ 2 & 3 & -6 \\ -11 & 2 & 0 \end{array}$$

$$\frac{x_1}{37} = \frac{x_2}{12} = \frac{x_3}{66} \quad X_3 = \begin{pmatrix} 37 \\ 12 \\ 66 \end{pmatrix}$$

Note:- To find the eigen vectors corresponding to nonrepeated eigen value of a matrix, we proceed as follows:-

- 1) Select the 1st two rows only
- 2) Start from the 1st row middle no. & move in anticlockwise direction to complete 1 cycle. If any element exist in the main diagonal while we are moving in anticlockwise direction, then the eigen value should be subtracted from the corresponding diagonal elements.

These elements will be return separately in row wise

- 3) Repeat the same procedure with 2nd row also.
& find eigen vectors

* Cayley - Hamilton theorem *

Statement:

Every square matrix satisfies its own characteristic eqⁿ.

Ex. Consider the matrix $A = \begin{pmatrix} 4 & 2 \\ 2 & 4 \end{pmatrix}$

Its char. eqⁿ is $\lambda^2 - 8\lambda + 12 = 0$

by Cayley - Hamilton theorem, every square matrix satisfies its own char. eqⁿ.

$$\text{i.e., } A^2 - 8A + 12I = 0$$

Note: (2x2)

Consider the matrix $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$

$$|A - \lambda I| = 0$$

$$\begin{vmatrix} a-\lambda & b \\ c & d-\lambda \end{vmatrix} = 0$$

$$(a-\lambda)(d-\lambda) - bc = 0$$

$$ad - a\lambda - d\lambda + \lambda^2 - bc = 0$$

$$\lambda^2 - (a+d)\lambda + (ad-bc) = 0$$

$$\Rightarrow \lambda^2 - \lambda[\text{trace of } A] + |A| = 0$$

by Cayley-Hamilton theorem, it is

$$A^2 - A[\text{trace of } A] + |A| \cdot I = 0$$

chc

cay

Ha

Note: (3×3)

Consider the 3×3 matrix

$$A = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}, \text{ its characteristic eq. is -}$$

General procedure: -

$$\lambda^3 - \lambda^2 (\text{trace of } A) + \lambda \{ | \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix} | + | \begin{vmatrix} a_{11} & a_{13} \\ a_{31} & a_{33} \end{vmatrix} | + | \begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix} | \} - |A| = 0$$

$$-|A| = 0$$

$$\begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix}$$

$$\begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix}$$

$$\begin{vmatrix} a_{11} & a_{13} \\ a_{31} & a_{33} \end{vmatrix}$$

Using Cayley Hamilton theorem, we can find

→ Inverse of a matrix

→ Powers of a matrix.

* Positive powers of matrix

$$A = \begin{pmatrix} 4 & 2 \\ 2 & 4 \end{pmatrix}$$

char. eqⁿ → $\lambda^2 - 8\lambda + 12 = 0$

Cayley Hamilton → $A^2 - 8A + 12I = 0$ ——— (1)

Hamilton
theorem

$$A^2 = 8A - 12I \quad \text{———— (2)}$$

$$A^3 = 8A^2 - 12A \quad \text{———— (3)}$$

$$A^4 = 8A^3 - 12A^2 \quad \text{———— (4)}$$

$$A^5 = 8A^4 - 12A^3 \quad \text{———— (5)}$$

⋮

* Negative powers of matrix

$$A^2 - 8A + 12I = 0$$

$$A^2 \cdot A^{-1} - 8A \cdot A^{-1} + 12I \cdot A^{-1} = 0$$

$$A - 8I + 12A^{-1} = 0$$

$$12A^{-1} = 8I - A$$

$$A^{-1} = \frac{1}{12} [-A + 8I]$$

$$A^{-1} \cdot A^{-1} = \frac{1}{12} [-\bar{A} \cdot A + 8I \bar{A}]$$

$$A^{-2} = \frac{1}{12} [-I + 8A^{-1}]$$

$$A^{-1} \cdot A^{-2} = \frac{1}{12} \left[-A^{-1} \cdot I + 8 A^{-1} \cdot A^{-1} \right]$$

$$A^{-3} = \frac{1}{12} \left[-A^{-1} + 8 A^{-2} \right]$$

Problem 1: — Find A^8 using Cayley hamilton theorem for

$$A = \begin{pmatrix} 1 & 2 \\ 2 & -1 \end{pmatrix}$$

$$|A| = -1 - 4 = -5$$

$$\text{trace of } A = 1 + (-1) = 0$$

$$A^2 - A(\text{trace of } A) + |A| \cdot I = 0$$

$$A^2 - A(0) + (-5) \cdot I = 0$$

$$A^2 + A - 5I$$

$$A^2 - 5I = 0$$

$$A^2 = 5I$$

$$(A^2)^4 = (5I)^4 = 5^4 \cdot I^4$$

$$A^8 = 625 I$$

$$= 625 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$A^8 = \begin{pmatrix} 625 & 0 \\ 0 & 625 \end{pmatrix}$$

(4m)
Problem 2:

Cayley-Hamilton theorem states that every square matrix satisfies its own characteristic eqⁿ.

Consider the matrix

$$A = \begin{pmatrix} -3 & 2 \\ -1 & 0 \end{pmatrix}$$

once we get char. eqⁿ after replacing λ by A

1st time we get char eqⁿ

we should not multiply it by A^T .

1) A satisfies the relation.

Ⓐ $A + 3I + 2A^T = 0$ Ⓑ $A^2 + 2A + 2I = 0$

Ⓒ $(A + I)(A + 2I) = 0$ Ⓓ $\exp(A)$

$$|A| = 0 + 2 = 2$$

$$\text{trace of } A = -3 + 0 = -3$$

$$\Rightarrow A^2 - A(\text{trace of } A) + |A| \cdot I = 0$$

$$A^2 + 3A + 2I = 0$$

$$A^2 + 2A$$

$$A^2 + 2AI + A + 2I = 0$$

$$A(A + 2I) + I(A + 2I) = 0$$

$$(A + 2I)(A + I) = 0$$

2) A^9 equals $511A + 510I$

a) $511A + 510I$ c) $154A + 155I$

b) $309A + 104I$ d) $\exp(9A)$

\Rightarrow

$$A^2 + 3A + 2I = 0$$

$$A^2 = -3A - 2I \quad \text{--- (1)}$$

$$A^3 = -3A^2 - 2A \quad \text{--- (2)}$$

$$(-3 > 2)$$

Not in
 series of
 ...

(even)
 $A \rightarrow$ all -ve elem
 $A \text{ odd} \rightarrow$ all +ve elem

$$\Rightarrow A^3 = -3(-3A - 2I) - 2A \quad (7 > 6)$$

$$A^3 = 7A + 6I \quad \text{--- (2)}$$

$$\Rightarrow A^4 = 7A^2 + 6A$$

$$= 7[-3A - 2I] + 6A$$

$$= -21A - 14I + 6A$$

$$A^4 = -15A - 14I \quad \text{--- (3)}$$

$$(15 > 14)$$

$$\Rightarrow A^5 = -15A^2 - 14A$$

$$= -15[-3A - 2I] - 14A$$

$$= 45A + 30I - 14A$$

$$A^5 = 31A + 30I \quad \text{--- (4)}$$

$$(31 > 30)$$

Imp *

$A^{\text{(Higher Number)}}$ $\Rightarrow A^{5000}, A^{10000}$

* Method :-

char eqⁿ $\Rightarrow \lambda^2 - (-3+0)\lambda + 2 = 0$

$$\lambda^2 + 3\lambda + 2 = 0$$

$$\lambda = -1, -2$$

Pro

\Rightarrow If λ values are not repeated
 then

$$\lambda^n = a\lambda + b \quad \text{--- (1)}$$

for $\lambda = -1$ $(-1)^n = -a + b \quad \text{--- (2)}$

for $\lambda = -2$ $(-2)^n = -2a + b \quad \text{--- (3)}$

$$(-1)^n - (-2)^n a = a$$

$$\therefore \boxed{a = \frac{(-1)^n - (-2)^n}{-1 - (-2)}}$$

$$\therefore \boxed{b = (-1)^n + a}$$

$$b = (-1)^n + a$$

$$= (-1)^n + (-1)^n - (-2)^n$$

$$= 2(-1)^n - (-2)^n$$

put a & b in eqⁿ. (1)

$$\lambda^n = [(-1)^n - (-2)^n] \lambda + [2(-1)^n - (-2)^n]$$

by c-b theorem

$$A^n = [(-1)^n - (-2)^n] A + [2(-1)^n - (-2)^n] I$$

For ex. put $n = 9$

$$A^9 = [(-1)^9 - (-2)^9] A + [2(-1)^9 - (-2)^9] I$$

$$= (-1 + 512) A + (-2 + 512) I$$

$$A^9 = 511 A + 510 I$$

Problem 3: The char eqⁿ. of 3×3 matrix P is given by

$$q(\lambda) = |\lambda I - P| = \lambda^3 + \lambda^2 + 2\lambda + 1$$

where I denoted the Identity matrix. The inverse of the matrix P will be —

a) $P^2 + P + 2I$

c) $-(P^2 + P + I)$

b) $P^2 + P + I$

d) $-(P^2 + 2P + 2I)$

$$\Rightarrow P^3 + P^2 + 2P + I = 0$$

$$-P = -I - P^2 - P^3$$

$$P^{-1} \cdot P^3 + P^{-1} \cdot P^2 + 2P^{-1} P + P^{-1} I = 0$$

$$P^2 + P + 2I + P^{-1} = 0$$

$$P^{-1} = -P^2 - P - 2I = -(P^2 + P + 2I)$$

GradeProblem (E. value & E. vectors)

How many L.I. eigen vectors of $A = \begin{pmatrix} 2 & 1 \\ 0 & 2 \end{pmatrix}$

$$\lambda = 2, 2$$

No. of L.I. e. vectors $\Rightarrow 1 \leq p \leq \textcircled{m}$ ← No. of times
 \downarrow e. value
 2 repeated

$$p = n - r$$

no of unknowns

$$r(A - \lambda I)$$

$$(A - \lambda I) = \begin{pmatrix} 2 - \lambda & 1 \\ 0 & 2 - \lambda \end{pmatrix} \quad \lambda = 2$$

$$= \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$$

$$r(A - \lambda I) = 1$$

$$p = n - r$$

$$= 2 - 1 = \boxed{1}$$

————— \times —————